

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 May 2007 (18.05.2007)

PCT

(10) International Publication Number
WO 2007/053946 A1

(51) International Patent Classification:

A61K 47/48 (2006.01) A61P 1/08 (2006.01)
A61K 38/26 (2006.01) A61P 3/04 (2006.01)
A61K 47/42 (2006.01) A61P 3/10 (2006.01)

(21) International Application Number:

PCT/CA2006/001835

(22) International Filing Date:

9 November 2006 (09.11.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/735,404 9 November 2005 (09.11.2005) US

(71) Applicant (for all designated States except US): CON-
JUCHEM BIOTECHNOLOGIES INC. [CA/CA]; 225,
Président-Kennedy, Third Floor, Suite 3950, Montréal,
Quebec H2X 3Y8 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BRIDON, Do-
minique, P. [FR/US]; 2167, Bay Street, San Francisco,
CA 94123 (US). CASTAIGNE, Jean-Paul [CA/CA]; 455
Lockhart Avenue, Mount-Royal, Quebec H3P 1Y6 (CA).
THIBODEAU, Karen [CA/CA]; 700, des Vignobles,
Rosemeres, Quebec J7A 4T2 (CA).

(74) Agent: BERESKIN & PARR; 40 King Street West, 40th
Floor, Suite 4000, Toronto, Ontario, M5H 3Y2 (CA).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS,
LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY,
MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS,
RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF TREATING DIABETES AND/OR OBESITY WITH REDUCED NAUSEA SIDE EFFECTS USING
AN INSULINOTROPIC PEPTIDE CONJUGATED TO ALBUMIN

(57) Abstract: The present invention provides methods of administering an insulinotropic peptide in an amount effective to treat
a disorder or condition while reducing nausea side effect by administering to a subject in need thereof an insulinotropic peptide
conjugated to albumin. The present invention also provides methods of selecting a subject for administration of a conjugated in-
sulinotropic peptide. Exemplary disorders or conditions treatable with an insulinotropic peptide include obesity and type II diabetes.

**METHOD OF TREATING DIABETES AND/OR OBESITY WITH REDUCED NAUSEA SIDE EFFECTS
USING AN INSULINOTROPIC PEPTIDE CONJUGATED TO ALBUMIN**

[0001] This application claims benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 60/735,404, filed November 9, 2005, which is hereby incorporated by reference in its entirety.

1. FIELD OF THE INVENTION

[0002] The present invention provides methods of administering an insulinotropic peptide in an amount effective to treat a disorder or condition while reducing nausea. Exemplary disorders or conditions treatable with an insulinotropic peptide include obesity and type II diabetes.

2. BACKGROUND OF THE INVENTION

[0003] The prevalence of diabetes for all age groups worldwide was estimated to be 2.8%, or 171 million in 2000, and is projected to be 4.4%, or 366 million in 2030. *See Wild et al., 2004, Diabetes Care 27(5):1047-1053.* In the United States alone, the prevalence of diabetes mellitus in 2005 was estimated at 20.8 million, or roughly 7% of the U.S. population. *See Centers for Disease Control and Prevention, 2005, National Diabetes Fact Sheet: General Information and National Estimates on Diabetes in the United States, 2005.* Approximately 95% of all patients with diabetes mellitus have type II disease. Diabetes is currently the fifth leading cause of death in the United States and is associated with excess morbidity stemming from cardiovascular disease, kidney failure, blindness, and lower limb amputation.

[0004] Similarly, obesity is a condition increasingly affecting the population worldwide. According to the World Health Organization, in 1995 there were an estimated 200 million obese adults worldwide and another 18 million under-five children classified as overweight. As of 2000, the number of obese adults had increased to over 300 million. *See Formiguera et al., 2004, Best Practice & Research Clinical Gastroenterology, 18:6, 1125-1146.*

[0005] The insulinotropic peptides glucagon-like peptide (GLP-1), exendin-3 and exendin-4 have been investigated as possible therapeutic agents for the management of type II non-insulin-dependent diabetes mellitus as well as related metabolic disorders, such as obesity. GLP-1 is a proglucagon-derived peptide secreted from intestinal L-cells in response to nutrient ingestion. GLP-1 acts as an incretin to lower postprandial glycemic excursion via

stimulation of insulin secretion and inhibition of glucagon secretion. GLP-1 also exerts actions independent of islet hormone secretion, including inhibition of both gastric emptying and food intake and stimulation of β -cell proliferation. As GLP-1 agonists, exendin-3 and exendin-4 mimic the actions of naturally occurring GLP-1 and are therefore classified as incretin mimetics.

[0006] While useful, GLP-1, exendin-3 and exendin-4 can present limited duration of action associated with short plasma half-lives *in vivo*, mainly due to rapid serum clearance and proteolytic degradation by dipeptidyl peptidase-IV. In addition, administration of these peptides for the treatment of diabetes or obesity can result in nausea side effect.

Subcutaneous injection of GLP-1 at high doses (1.5 nmol/kg body weight and higher) lead to nausea and, less often, vomiting in a significant proportion of subjects exposed to treatment. See Ritzel *et al.*, 1995, *Diabetologia* 38:720-725. Further, in three phase III comparative efficacy trials of exenatide, the synthetically derived peptide of exendin-4, nausea was the most commonly reported adverse event, with an incidence of 43.5% of the 1446 enrolled patients enrolled in all three studies. See Iltz *et al.*, 2006, *Clin. Ther.* 28(5): 652-665.

[0007] Thus, there is a need for administration of forms of GLP-1, exendin-3, exendin-4, and other insulintropic peptides which minimize side effects such as nausea and vomiting, but maintain their therapeutic advantages.

3. SUMMARY OF THE INVENTION

[0008] The present invention is based, in part, on the discovery that administration to a subject of an insulintropic peptide conjugated to albumin can prevent, reduce, or eliminate nausea side effect compared to administration of the insulintropic peptide alone.

Accordingly, the present invention provides methods of administering to a subject a form of an insulintropic peptide that prevents, reduces, or eliminates nausea side effect compared to administration of the insulintropic peptide alone. In certain embodiments the insulintropic peptide is in the form of a conjugate. In some embodiments the insulintropic peptide is in the form of an insulintropic peptide conjugated to a macromolecule. In some embodiments the insulintropic peptide is in the form of an insulintropic peptide conjugated to albumin.

[0009] In certain embodiments, the methods of the invention comprise the step of administering to the subject an insulintropic peptide conjugated to albumin that can prevent, reduce, or eliminate nausea side effect that may accompany administration of the unconjugated insulintropic peptide. In some embodiments, administration of the conjugated insulintropic peptide prevents nausea side effect compared to administration of the

unconjugated insulinotropic peptide. In some embodiments, administration of the conjugated insulinotropic peptide reduces nausea side effect compared to administration of the unconjugated insulinotropic peptide. In some embodiments, administration of the conjugated insulinotropic peptide eliminates nausea side effect compared to administration of the unconjugated insulinotropic peptide. In some embodiments, the amount administered is effective to prevent, reduce, or eliminate nausea side effect in the subject.

[0010] While not intending to be bound by any particular theory of operation, it is believed that an effective approach to prevent, reduce, or eliminate the incidence of nausea that can accompany the administration of an insulinotropic peptide is to identify a patient population having a sensitivity to nausea before, during or after treatment with an insulinotropic peptide. Accordingly, the patient population can be treated with reduced nausea by administration of an insulinotropic peptide conjugated to albumin as described herein.

[0011] In certain embodiments, the present invention provides methods of selecting a population, for example a sub-population, of subjects suitable for treatment with the insulinotropic peptide conjugated to albumin. Exemplary methods of selection are described herein.

[0012] In certain embodiments, the methods of the invention comprise the step of administering to the subject an amount of conjugated insulinotropic peptide effective for treating a disorder or condition while providing reduced nausea side effect. Exemplary conditions or disorders treatable with insulinotropic peptides include, but are not limited to, obesity and diabetes, including maturity onset diabetes mellitus (type II diabetes).

[0013] The insulinotropic peptide of the present invention can be any insulinotropic peptide known to those of skill in the art. For example it can be any peptide which stimulates, or causes the stimulation of, the synthesis or expression of the hormone insulin. In some embodiments the insulinotropic peptide is selected from the group consisting of glucagon-like peptide 1, exendin-3 and exendin-4, and their precursors, derivatives, or fragments.

[0014] The conjugated insulinotropic peptide of the present invention can be conjugated to a macromolecule. In some embodiments, the insulinotropic peptide is conjugated to albumin. In some embodiments, the insulinotropic peptide is conjugated to serum albumin. In some embodiments, the insulinotropic peptide is conjugated to human serum albumin. In some embodiments, the insulinotropic peptide is conjugated to

recombinant serum albumin. In some embodiments, the insulinotropic peptide is conjugated to recombinant human serum albumin.

[0015] The conjugated insulinotropic peptide of the present invention can be administered in a pharmaceutically acceptable formulation by any route known to those of skill in the art, including but not limited to oral, intranasal, intravascular, intraarterial, intramuscular, subcutaneous, transdermal or rectal administration.

[0016] In another aspect, the present invention provides methods of selecting a subject for treatment with an insulinotropic peptide conjugated to albumin. In certain embodiments, the methods comprise the step of identifying a subject having a condition or disorder treatable with an insulinotropic peptide who is prone to experiencing, has experienced, or is experiencing nausea. The subject is selected for treatment when the subject is prone to experiencing, has experienced, or is experiencing nausea as determined by any method known to those of skill in the art without limitation. Exemplary methods are described herein.

[0017] In further aspect, the present invention provides methods of administering an insulinotropic peptide. In certain embodiments, the methods comprise the step of selecting a nausea-sensitive subject having a disorder or condition treatable with the insulinotropic peptide and administering to the subject the insulinotropic peptide conjugated to albumin in an amount effective to treat the disorder or condition, wherein the conjugated insulinotropic peptide prevents, reduces, or eliminates nausea compared to the administration of unconjugated insulinotropic peptide. The methods of selecting a subject who is prone to experiencing has experienced or is experiencing nausea are described herein.

4. BRIEF DESCRIPTION OF THE FIGURES

[0018] FIG. 1 presents a graph representing mean (\pm SEM) saccharin preference ratios in rats following intraperitoneal and subcutaneous administration of unconjugated exendin-4(1-39) versus exendin-4(1-39) Lys⁴⁰ ϵ -AEEA-MPA conjugated with albumin (CJC-1134-PC), 72-84 hours post dose;

[0019] FIG. 2 presents a graph representing mean (\pm SEM) saccharin preference ratios in rats following intraperitoneal and subcutaneous administration of unconjugated exendin-4(1-39) versus CJC-1134-PC, 96-108 hours post dose;

[0020] FIG. 3 presents a graph representing mean cumulative kaolin intake (g/animal) in rats following administration of unconjugated exendin-4(1-39) versus CJC-1134-PC, 0-8 hours post dose;

- [0021] FIG. 4 presents a graph representing mean (\pm SEM) food consumption (g/animal/day) over 7 days in rats following administration of CJC-1134-PC (0, 1, 5, and 25 mg/kg);
- [0022] FIG. 5 presents a graph representing mean (\pm SEM) body weight (g) over 7 days in rats following administration of CJC-1134-PC (0, 1, 5, and 25 mg/kg);
- [0023] FIG. 6 presents a graph representing mean (\pm SEM) food consumption (g/animal/day) over 28 days in rats (n=15) following administration of CJC-1134-PC (25 mg/kg) and over a two-week recovery (n=5);
- [0024] FIG. 7 presents a graph representing delta (\pm SEM) body weight change (g) over 28 days in rats (n=15) following administration of CJC-1134-PC (25 mg/kg) and over a two week recovery (n=5);
- [0025] FIG. 8 presents a graph representing delta (\pm SEM) food consumption (number of biscuits) over 7 days in monkeys following administration of CJC-1134-PC (0, 0.4, 2, and 10 mg/kg);
- [0026] FIG. 9 presents a graph representing delta (\pm SEM) body weight change (kg) over 7 days in monkeys following administration of CJC-1134-PC (0, 0.4, 2, and 10 mg/kg);
- [0027] FIG. 10 presents a graph representing mean (\pm SEM) food consumption (number of biscuits) over 28 days in monkeys (n=6) following administration of CJC-1134-PC (10 mg/kg) and over a two-week recovery (n=3);
- [0028] FIG. 11 presents a graph representing delta (\pm SEM) body weight change (kg) over 28 days in monkeys (n=6) following administration of CJC-1134-PC (10 mg/kg) and over a two-week recovery (n=3);
- [0029] FIG. 12 presents a graph representing blood glucose concentrations (nmol/L) over 24 hours in female db/db mice following a single subcutaneous administration of unconjugated exendin-4(1-39) versus CJC-1134-PC;
- [0030] FIG. 13 presents a graph representing blood glucose concentrations (nmol/L) over 48 hours in female db/db mice following administration of unconjugated exendin-4(1-39) versus CJC-1134-PC;
- [0031] FIG. 14 presents a graph representing blood glucose concentrations (nmol/L) over 101 hours in female db/db mice following administration of unconjugated exendin-4(1-39) versus CJC-1134-PC;
- [0032] FIG. 15 presents a graph representing food consumption (g/animal) over 101 hours in female db/db mice following administration of unconjugated exendin-4(1-39) versus CJC-1134-PC;

[0033] FIG. 16 presents a graph representing water consumption (g/animal) over 101 hours in female db/db mice following administration of unconjugated exendin-4(1-39) versus CJC-1134-PC;

[0034] FIG. 17 presents a graph representing mean (\pm SEM) cumulative weight gain (%) over 25 days in male Zucker Diabetic Fatty (ZDF) rats following administration of unconjugated exendin-4(1-39) versus CJC-1134-PC; and

[0035] FIG. 18 presents a graph representing mean (\pm SEM) cumulative food intake (g/animal) over 25 days in male ZDF rats following administration of unconjugated exendin-4(1-39) versus CJC-1134-PC.

5. DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5.1 Definitions

[0036] As used herein, the following terms shall have the following meanings unless otherwise specified:

[0037] "Subject" refers to an animal such as a mammal, including but not limited to, a primate (e.g., human), cow, sheep, goat, horse, dog, cat, rabbit, rat, mouse and the like. In preferred embodiments, the subject is human. In certain embodiments, the subject is a non-human animal, for instance, a non-human animal such as a cow, sheep, goat, horse, dog, cat, rabbit, rat or mouse.

[0038] "Nausea" refers to any unpleasant sensation in the epigastrium, in the back of the throat, or in the abdomen. In some embodiments, nausea may culminate in vomiting. In some embodiments, nausea may not culminate in vomiting. In some embodiments, nausea can be sickness at the stomach, especially when accompanied by a loathing for food and an involuntary impulse to vomit. In some embodiments, nausea can be a feeling of sickness or discomfort in the stomach marked by an urge to vomit.

[0039] A "nausea-sensitive subject" is any subject who is prone to experiencing nausea in response to certain physical, chemical, and / or psychological stimuli. Exemplary stimuli are described herein. In some embodiments, the subject has a medical history of nausea sensitivity. In some embodiments, the subject has recently experienced nausea independent of administration of an insulinotropic peptide. In some embodiments, the subject has recently experienced nausea in response to administration of an insulinotropic peptide. In some embodiments, the subject is currently experiencing nausea independent of administration of an insulinotropic peptide. In some embodiments, the subject is currently experiencing nausea in response to administration of an insulinotropic peptide.

[0040] "Reducing," "reduces" or "reduced nausea side effect" refers to a reduction in the degree, duration, and / or frequency of nausea experienced by a subject following administration of an insulinotropic peptide conjugated to albumin compared to administration of an unconjugated insulinotropic peptide. Such reduction encompasses the prevention of nausea that a subject would otherwise experience in response to an unconjugated insulinotropic peptide. Such reduction also encompasses the elimination of nausea side effect previously experienced by a subject following administration of an unconjugated insulinotropic peptide. In some embodiments, "reducing" or "reduced nausea" encompasses a reduction of nausea wherein the nausea is reduced to zero or undetectable levels. In other embodiments, nausea is reduced but not completely eliminated.

[0041] As used herein, "compared to unconjugated insulinotropic peptide" is used interchangeably with "compared to insulinotropic peptide alone" and refers to amounts known to be comparable according to one of skill in the art. In some embodiments, the comparable amounts are equimolar amounts. In some embodiments, the comparable amounts are weight to weight equivalents.

[0042] "Insulinotropic peptide" refers to a peptide with insulinotropic activity. Insulinotropic peptides can stimulate, or cause the stimulation of, the synthesis or expression of the hormone insulin. Such peptides include precursors, derivatives, or fragments of peptides such as glucagon-like peptide, exendin-3 and exendin-4 and other peptides with insulinotropic activity.

[0043] "Glucagon-Like Peptide" ("GLP") and "GLP derivatives" are intestinal hormones which generally simulate insulin secretion during hyperglycemia, suppress glucagon secretion, stimulate (pro) insulin biosynthesis and decelerate gastric emptying and acid secretion. In some embodiments, the glucagon-like peptide is GLP-1(7-37). In some embodiments, the glucagon-like peptide is GLP-1(7-36). Some GLPs and GLP derivatives promote glucose uptake by cells but do not simulate insulin expression, as disclosed in U.S. Pat. No. 5,574,008 which is hereby incorporated by reference.

[0044] "Exendin-3" is a naturally occurring GLP-1 agonist isolated from salivary secretions of *Heloderma horridum*, the Mexican bearded lizard, and shares a 53% overlap with mammalian GLP-1 amino acid sequence, as disclosed in U.S. Pat. No. 5,424,286 which is hereby incorporated by reference.

[0045] "Exendin-4" is a naturally occurring GLP-1 agonist isolated from salivary gland venom of *Heloderma suspectum*, the Gila monster, and shares a 53% overlap with mammalian GLP-1 amino acid sequence as disclosed in U.S. Pat. No. 5,424,286 which is

hereby incorporated by reference. Exendin-4 decreases glucagons and increases insulin secretion in a glucose-dependent manner, and mimics certain actions of GLP-1, including binding to and activating the human GLP-1 receptor. Exendin-4 improves glycemic control by reducing fasting and postprandial glucose concentrations through restoration of first-phase insulin response, regulation of glucagon secretion, delaying gastric emptying, and decreasing food intake.

[0046] "Reactive groups" are chemical groups capable of forming a covalent bond. Such reactive agents can be coupled or bonded to an insulintropic peptide of interest to form a modified insulintropic peptide. Reactive groups can generally be carboxy, phosphoryl, or convenient acyl group, either as an ester or a mixed anhydride, or an imidate, thereby capable of forming a covalent bond with functionalities such as an amino group, a hydroxy group or a thiol group at the target site on albumin. For the most part, the esters will involve phenolic compounds, or be thiol esters, alkyl esters, phosphate esters, or the like. Reactive groups include succinimidyl-containing groups and maleimido-containing groups.

[0047] "Functionalities" are groups on albumin to which reactive groups on modified insulintropic peptides are capable of reacting with to form covalent bonds. Functionalities include hydroxyl groups for bonding to ester reactive entities; thiol groups for bonding to maleimides and maleimido-containing groups, imidates and thioester groups; and amino groups for bonding to carboxy groups, phosphoryl groups, acyl groups, succinimidyl-containing groups or maleimido-containing groups on reactive entities.

[0048] "Linking Groups" are chemical moieties capable of connecting reactive groups to insulintropic peptides. Linking groups may comprise one or more alkyl groups such as methyl, ethyl, propyl, butyl, etc. groups, alkoxy groups, alkenyl groups, alkynyl groups or amino group substituted by alkyl groups, cycloalkyl groups, polycyclic groups, aryl groups, polyaryl groups, substituted aryl groups, heterocyclic groups, and substituted heterocyclic groups. Linking groups may also comprise poly ethoxy aminoacids such as AEA ((2-amino) ethoxy acetic acid) or a preferred linking group AEEA ([2-(2-amino)ethoxy])ethoxy acetic acid). In certain embodiments, a linking group can comprise, for example, a chain of 0-30 atoms, or 0-20 atoms, or 0-10 atoms. In certain embodiments, a linking group can consist of, for example, a chain of 0-30 atoms, or 0-20 atoms, or 0-10 atoms. Those atoms can be selected from the group consisting of, for example, C, N, O, S, P.

[0049] As used herein, "albumin" refers to the most abundant protein in blood plasma having a molecular weight of approximately between 65 and 67 kilodaltons in its monomeric form, depending on the species of origin. The term "albumin" is used interchangeably with

“serum albumin” and is not meant to define the source of the albumin which forms a conjugate with the modified peptides of the invention. Thus, the term “albumin” as used herein may refer either to albumin purified from a natural source such as blood or serous fluids, or it may refer to chemically synthesized or recombinantly produced albumin.

[0050] A “conjugated insulintropic peptide” is an insulintropic peptide that has been conjugated to macromolecule, for example albumin, via a covalent bond formed between the insulintropic peptide and a functionality on the macromolecule. In some embodiments, the insulintropic peptide has been modified to contain a reactive group to which the macromolecule is covalently bound. In some embodiments, the reactive group is coupled to the insulintropic peptide via a linking group.

[0051] “Preventing” or “prevention” of any disease or disorder refers to a reduction in the risk of acquiring a disease or disorder (*i.e.*, causing at least one of the clinical symptoms of the disease not to develop in a subject that may be exposed or predisposed to the disease but does not yet experience or display symptoms of the disease). Preferably, prevention refers to the use of a compound or composition in a subject not yet affected by the disease or disorder or not yet exhibiting a symptom of the disease or disorder, for instance a subject not yet infected or not yet exhibiting the symptoms of infection.

[0052] “Treating” or “treatment” of any disease or disorder refers, in one embodiment, to ameliorating the disease or disorder (*i.e.*, arresting or reducing the development of the disease or at least one of the clinical symptoms thereof) that exists in a subject. In another embodiment, “treating” or “treatment” refers to ameliorating at least one physical parameter, which may be indiscernible by the subject. In yet another embodiment, “treating or treatment” refers to modulating the disease, either physically (*e.g.*, stabilization of a discernable symptom) or physiologically (*e.g.*, stabilization of a physical parameter) or both. In yet another embodiment, “treating” or “treatment” refers to delaying the onset of the disease or disorder.

[0053] Effective amount: “an effective amount” means an amount of a conjugated insulintropic peptide that when, administered to a subject for treating a disease is sufficient to effect such treatment for the disease. An effective amount can vary depending on, inter alia, the insulintropic peptide used, the disease and its severity and the age, weight, etc. of the subject to be tested. In certain embodiments, the amount is effective to treat diabetes or obesity. In certain embodiments, the amount is effective to treat type 1 diabetes or type 2 diabetes. In certain embodiments, the amount is effective to lower blood glucose, to lower whole blood glucose, to lower plasma blood glucose and/or to lower HbA1c levels. In

certain embodiments, the levels can be lowered by 5%, 10%, 15% or more. In certain embodiments, the levels on average can be lowered for at least one day, at least two days, at least three days, at least four days or least one week.

5.2 Methods of Treatment with Reduced Nausea

[0054] Although not intending to be bound by any particular theory of operation, the present invention is based, in part, on the discovery that certain populations of subjects receiving insulinotropic peptides for the treatment of a disorder or condition experience nausea side effect in response to the administration of the insulinotropic peptide. It is further discovered that administration to a subject of an insulinotropic peptide conjugated to albumin can reduce nausea side effect compared to administration of the insulinotropic peptide alone.

[0055] Accordingly, the present invention provides methods of treating a disorder or condition with a form of insulinotropic peptide that provides reduced nausea. The methods comprise the step of administering to the subject an amount of an insulinotropic peptide conjugated to albumin effective to treat the disorder or condition with reduced nausea side effect.

[0056] Exemplary disorders or conditions treatable with an insulinotropic peptide include, but are not limited to, obesity and diabetes, including maturity onset diabetes mellitus (type II diabetes). In certain embodiments, the disorder or condition treatable with an insulinotropic peptide is obesity. In another embodiment, the disorder or condition treatable with an insulinotropic peptide is diabetes. In certain embodiments, the disorder or condition treatable with an insulinotropic peptide is type II diabetes.

[0057] Nausea of the present invention can be any unpleasant sensation in the epigastrium, in the back of the throat, or in the abdomen. In some embodiments, nausea may culminate in vomiting. In some embodiments, nausea may not culminate in vomiting. In some embodiments, nausea can be sickness at the stomach, especially when accompanied by a loathing for food and an involuntary impulse to vomit. In some embodiments, nausea can be a feeling of sickness or discomfort in the stomach marked by an urge to vomit. In some embodiments, the nausea is mild. In some embodiments, the nausea is moderate. In some embodiments, the nausea is severe. In some embodiments the nausea does not interfere with the normal daily life of the subject. In some embodiments the nausea interferes with normal daily life. In some embodiments the subject is bedridden because of the nausea.

[0058] In one aspect, the present invention provides methods of preventing nausea in a subject in response to the administration of an insulinotropic peptide. In another aspect, the present invention provides methods of reducing nausea in a subject in response to the

administration of an insulinotropic peptide. In another aspect, the present invention provides methods of eliminating nausea in a subject in response to the administration of an insulinotropic peptide.

[0059] It should be recognized that the methods of the invention encompass the prevention, reduction, and elimination of nausea in a subject receiving an insulinotropic peptide for the treatment of a disorder or condition by selecting patients who are prone to experiencing, have experienced, or are experiencing nausea. Those more likely to benefit from reduced nausea are identified by determining their nausea sensitivity either resulting from, or arising independent of, the administration of an unconjugated insulinotropic peptide.

5.2.1 Subjects

[0060] In certain embodiments of the invention, the subject is an animal, preferably a mammal, more preferably a non-human primate. In certain embodiments, the subject is a human patient. The subject can be a male or female subject. In certain embodiments, the subject is a non-human animal, such as, for instance, a cow, sheep, goat, horse, dog, cat, rabbit, rat or mouse.

[0061] The methods of the invention can be used for selecting a subject for treatment with an insulinotropic peptide conjugated to albumin. Particularly useful subjects include those that have a disorder or condition treatable with an insulinotropic peptide that are prone to experiencing, have experienced, or are experiencing nausea.

[0062] In certain embodiments, the subject is at risk for a disorder or a condition treatable with an insulinotropic peptide including, but not limited to, obesity and type II diabetes. In some embodiments the subject is at risk for obesity. In some embodiments the subject is at risk for type II diabetes.

[0063] In some embodiments, the subject is not healthy. In some embodiments the subject has or suffers from a condition treatable with an insulinotropic peptide including, but not limited to, obesity or type II diabetes.

[0064] In some embodiments, the subject is obese. In some embodiments, the subject is a human and has a Body Mass Index (BMI) of 30 kg/m^2 or greater. In some embodiments, the subject is a human and has a BMI between 30 kg/m^2 and 35 kg/m^2 . In some embodiments, the subject is a human and has a BMI of 35 kg/m^2 or greater. In some embodiments, the subject is a human and has a BMI of 40 kg/m^2 or greater. In some embodiments, the subject weighs more than 120% of the normal weight for its age and height. In some embodiments, the subject is a human and weighs more than 96 kg.

[0065] In some embodiments, the subject has type II diabetes. In some embodiments, the subject has abnormal glucose levels. In particular embodiments, the subject has a high glucose level. In some embodiments, the subject is a human and has an average whole blood glucose level of 8 mmol/L (138 mg/dl) or greater, and an average plasma blood glucose level of 9.0 mmol/L (154 mg/dl) or greater. In some embodiments, the subject is a human and has an average whole blood glucose level between 8 mmol/L (138 mg/dl) and 16 mmol/L (281 mg/dl), and an average plasma blood glucose level between 9.0 mmol/L (154 mg/dl) and 17 mmol/L (314 mg/dl). In some embodiments, the subject is a human and has an average whole blood glucose level greater than 16 mmol/L (281 mg/dl), and an average plasma blood glucose level greater than 17 mmol/L (314 mg/dl).

[0066] In some embodiments, the subject is a human and has a glycosylated hemoglobin (HbA1c) level of 6.5% or greater. In some embodiments, the subject is a human and has a HbA1c level between 6.5% and 11%. In some embodiments, the subject is human and has a HbA1c level of 11% or greater.

5.2.2 Assessing Nausea in Subjects Having a Disorder or Condition Treatable with an Insulinotropic Peptide

[0067] Whether a subject is prone to experiencing, has experienced, or is experiencing nausea can be determined according to any technique known to those of skill in the art without limitation. In certain embodiments, the technique for assessing nausea is not critical for the invention. In certain embodiments, the assessment of nausea is determined according to the judgment of the practitioner in the art. In certain embodiments, the assessment of nausea is determined by the subjective experience of the subject. In certain embodiments nausea is determined by the subjective experience of the subject combined with objective elements such as pallor, sweating and feeling cold.

[0068] In some embodiments the assessment of nausea is determined according to the Duke Descriptive Scale (DDS). See Laszlo *et al.*, 1981, *J. Clin. Pharmacol.* 2:51S-56S; Cotanch, 1983, *Cancer Nurs.* 6:277-283; the entirety of which are incorporated by reference herein. The DDS grades nausea on a scale of I to IV, taking into account intensity, severity, and impairment in patient activity for a 24-hour period. In grade I, the subject has experienced no nausea. In grade II, the nausea is mild, with no interference in daily activities. In grade III, the nausea is moderate, with some interference in daily activities. In grade IV, the nausea is severe, and the subject is bedridden with nausea for more than 2 hours.

[0069] In some embodiments the assessment of nausea is determined according to the Visual Analog Scale (VAS). See Bennett *et al.*, 1989, *Oncol. Nurs. Forum* 16:175, the entirety of which is incorporated by reference herein. The VAS is a line, usually 100 mm in length, occasionally 150 or 160 mm long, with anchors at each end to indicate the extremes of the nausea experienced by the subject. The low endpoint is to the left in a horizontally oriented scale, and at the base of a vertically oriented scale. Subjects indicate the point on the scale corresponding to the degree of nausea they are experiencing. Investigators score the intensity of the discomfort by measuring the millimeters from the low end of the scale to the mark.

[0070] In some embodiments the assessment of nausea is determined according to the Rhodes Index of Nausea and Vomiting (INV). See Rhodes *et al.*, 1994, *Cancer Nurs.* 17:45-51; Rhodes *et al.*, 1986, *Oncol. Nurs. Forum* 13:41-47; Rhodes *et al.*, 1988, *Appl. Nurs. Res.* 1:143-144; the entireties of each of which are incorporated by reference herein. The INV measures the individual components of nausea, vomiting, retching, and associated distress. This 8-item, 5-point scale measures a subject's perceived (1) duration of nausea; (2) frequency of nausea; (3) distress from nausea; (4) frequency of vomiting; (5) amount of vomiting; (6) distress from vomiting; and (7) frequency of retching. An exemplary INV item is: "During the last 12 hours, I have not felt any distress from nausea / sickness at my stomach"; "During the last 12 hours I have felt mild distress from nausea or sickness at my stomach"; "During the last 12 hours I have felt moderate distress from nausea or sickness at my stomach"; "During the last 12 hours I have felt great distress from nausea or sickness at my stomach"; "During the last 12 hours I have felt as severe distress from nausea or sickness at my stomach as can be."

[0071] In some embodiments the assessment of nausea is determined according to the Functional Living Index-Emesis (FLIE). See Lindley *et al.*, 1992, *Qual. Life Res.* 1:331-340, the entirety of which is incorporated by reference herein. The FLIE is a self-administered questionnaire which focuses on the effects of nausea and vomiting on physical activities, social and emotional function, and the ability to enjoy food. An exemplary FLIE item is: "How much nausea have you had in the past 3 days?" Each item is answered in a range from 1 to 7, with 9 items for nausea and 9 items for vomiting. A total score is created by adding the responses to the 18 questions. The range of total scores possible is between 18 (all 1 responses on the scale) and 126 (all 7 responses on the scale). Lower scores indicate a more negative impact of nausea and vomiting; higher scores indicate a higher impact.

[0072] In some embodiments the assessment of nausea is determined according to a daily diary of the subject. *See* Baltzer *et al.*, 1994, *Cancer* 73:720-723; Buser *et al.*, 1993, *Ann. Oncol.* 4:475-479; Sung *et al.*, 1993, *J. Clin. Anesthes.* 5:22-29; the entireties of each of which are incorporated by reference herein. Diaries have been used for periods ranging from 24 hours to 15 days. This method requires subject self-report and has been correlated to other measures such as observation and the FLIE.

5.2.3 Selecting Subjects for Treatment with a Conjugated Insulinotropic Peptide

[0073] Subjects having a disorder or condition treatable with an insulinotropic peptide are selected for treatment with an insulinotropic peptide conjugated to albumin if they are prone to experiencing, have experienced, or are experiencing nausea. Whether a subject is prone to experiencing, has experienced, or is experiencing nausea can be determined according to the judgment of the practitioner of the art. Exemplary methods are described herein.

[0074] In some embodiments, nausea of the present invention can be of any degree, duration, or frequency experienced by the subject as determined by the practitioner of the art. In some embodiments, nausea can be of any degree, duration, or frequency as determined by the subjective experience of the subject. In some embodiments, nausea can be of any degree, duration, or frequency according to the DDS. In some embodiments, nausea can be of any degree, duration, or frequency according to the VAS. In some embodiments, nausea can be of any degree, duration, or frequency according to the INV. In some embodiments, nausea can be of any degree, duration, or frequency according to the FLIE. In some embodiments, nausea can be of any degree, duration, or frequency according to a daily diary of the subject.

[0075] In certain embodiments, subjects who are prone to experiencing nausea are selected for treatment. In some embodiments, the subject has a medical history of nausea sensitivity independent of the administration of an unconjugated insulinotropic peptide. A subject who is prone to experiencing nausea may experience nausea due to any cause known in the art. For example, a subject prone to nausea may experience nausea in response to any physical, chemical, and / or psychological stimuli.

[0076] Exemplary stimuli include, but are not limited to: psychological problems, including anxiety, stress and depression; severe pain or shock; unpleasant smells or pictures; disturbances in the inner ear in conditions such as altitude sickness; Meniere's disease; motion sickness and labyrinthitis; diseases of the internal organs (viscera); gastrointestinal disease such as gastritis, peptic ulcer, stomach cancer, gut obstruction, appendicitis and

mesenteric adenitis; disease of the liver or gallbladder such as cholecystitis, hepatitis and cirrhosis; pancreatitis; disease of the heart such as myocarditis, severe hypertension, heart attack (myocardial infarction) and congestive heart failure; disease of the brain such as migraine, stroke (cerebrovascular accident), brain tumors, cerebral hypoxia, and epilepsy; diseases of the urinary tract such as urinary tract infections and renal colic; chemicals and toxins; bacterial and viral infections, such as cholera and salmonella, that can cause either gastro-intestinal or systemic infections; parasitic infections, such as malaria; meningitis; sinusitis; endocrine problems, such as hyperparathyroidism, hyperthyroidism and diabetes mellitus; waste products that accumulate in the body with renal failure, and other electrolyte disturbances; certain drugs and hormones including digoxin, morphine, codeine, estrogens, iron preparation and aminophylline; morning sickness; haemolytic anaemia; porphyria; alcoholism; radiotherapy; self-induced vomiting in psychiatric conditions such as bulimia nervosa; congenital malformations of the gastro-intestinal tract such as pyloric stenosis and gastro-oesophageal reflux; and constipation.

[0077] In some embodiments, subjects who have recently experienced nausea are selected for treatment. In some embodiments, the nausea was experienced independent of administration of an unconjugated insulintropic peptide. In other embodiments, the nausea was experienced as a result of a single administration of an unconjugated insulintropic peptide. In other embodiments, the nausea was experienced as a result of more than one administration of an unconjugated insulintropic peptide.

[0078] In some embodiments, the subject experienced acute nausea as a result of administration of the unconjugated insulintropic peptide. In some embodiments, the nausea was experienced within 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 hour(s) of a single administration of the unconjugated insulintropic peptide. In some embodiments, the nausea was experienced within 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 hour(s) of more than one administration of the unconjugated insulintropic peptide.

[0079] In other embodiments, the subject experienced delayed nausea as a result of administration of the unconjugated insulintropic peptide. In some embodiments, the nausea was experienced within 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 day(s) of a single administration of the unconjugated insulintropic peptide. In some embodiments, the nausea was experienced within 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 day(s) of more than one administration of the unconjugated insulintropic peptide.

[0080] In some embodiments, subjects who are currently experiencing nausea are selected for treatment. In some embodiments, the subject is experiencing nausea independent of administration of an unconjugated insulinotropic peptide. In other embodiments, the subject is experiencing nausea as a result of a single administration of an unconjugated insulinotropic peptide. In other embodiments, the subject is experiencing nausea as a result of more than one administration of an unconjugated insulinotropic peptide.

[0081] In some embodiments, the subject is experiencing acute nausea as a result of administration of the unconjugated insulinotropic peptide. In some embodiments, the subject is experiencing nausea within 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 hour(s) of a single administration of an unconjugated insulinotropic peptide. In some embodiments, the subject is experiencing nausea within 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 hour(s) of more than one administration of an unconjugated insulinotropic peptide.

[0082] In other embodiments, the subject is experiencing delayed nausea as a result of administration of the unconjugated insulinotropic peptide. In some embodiments, the subject is experiencing nausea within 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 day(s) of a single administration of an unconjugated insulinotropic peptide. In some embodiments, the subject is experiencing nausea within 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 day(s) of more than one administration of an unconjugated insulinotropic peptide.

[0083] In some embodiments, subjects are selected for treatment with a conjugated insulinotropic peptide of the present invention after a first instance of nausea in response to administration of an unconjugated insulinotropic peptide. In some embodiments, subjects are selected for treatment with a conjugated insulinotropic peptide after more than one instance of nausea in response to administration of an unconjugated insulinotropic peptide.

[0084] In addition to the assessment of nausea, other parameters or variables can be used in combination with the assessment of nausea to select a subject for treatment with an insulinotropic peptide conjugated to a macromolecule, for example albumin. In some embodiments, average whole blood glucose levels of the subject are used. In some embodiments, average plasma blood glucose levels of the subject are used. In some embodiments, HbA1c levels of the subject are used. In other embodiments, the body weight or BMI of the subject is used to make the selection.

5.2.4 Administration of the Conjugated Insulinotropic Peptide

[0085] The conjugated insulinotropic peptide can be administered according to any technique deemed suitable by one of skill in the art. In some embodiments, the conjugated

insulinotropic peptides can be administered in a physiologically acceptable medium, *e.g.* deionized water, phosphate buffered saline (PBS), saline, aqueous ethanol or other alcohol, plasma, proteinaceous solutions, mannitol, aqueous glucose, alcohol, vegetable oil, or the like. Other additives which may be included include buffers, where the media are generally buffered at a pH in the range of about 4.0 to 8.0, preferably in the range of about 5.0 to 8.0 and more preferably in the range of about 5.0 to 7.2, where the buffer will generally range in concentration from about 50 to 250 mM, salt, where the concentration of salt will generally range from about 5 to 500 mM, physiologically acceptable stabilizers, and the like. The compositions may be lyophilized for convenient storage and transport.

[0086] The conjugated insulinotropic peptides can for the most part be administered orally, parenterally, such as intravascularly (IV), intraarterially (IA), intramuscularly (IM), subcutaneously (SC), rectally, intradermally, intranasally, by inhalation or the like. Administration may in appropriate situations be by transfusion. Usually a single injection will be employed although more than one injection may be used, if desired. The conjugated insulinotropic peptides may be administered by any convenient means, including pen, needle-free device, enema, suppository, patch, inhaler, metered dose inhaler, syringe, trocar, catheter, or the like. The particular manner of administration will vary depending upon the amount to be administered, whether a single bolus or continuous administration, or the like. Preferably, the administration will be intravascularly, where the site of introduction is not critical to this invention, preferably at a site where there is rapid blood flow, *e.g.*, intravenously, peripheral or central vein. Also preferably, the administration will be subcutaneously.

5.2.5 Dosage and Frequency of Administration

[0087] The dosage and frequency of administration of the conjugated insulinotropic peptide will be determined according to one of ordinary skill in the art. The amount of a conjugated insulinotropic peptide in the methods of the invention which will be effective in the treatment of a disorder or condition with reduced nausea will vary with the nature and severity of the disorder or condition, and the route by which the active ingredient is administered. The frequency and dosage will also vary according to factors specific for each patient depending on the severity of the disorder or condition, the route of administration, as well as age, body weight, response, and the past medical history of the patient. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0088] Exemplary doses of a conjugated insulinotropic peptide include milligram or microgram amounts of the conjugated insulinotropic peptide per kilogram of subject or sample weight (e.g., about 1 mg to about 10 mg per dose, for example 1 mg, 3 mg, 5 mg or 10 mg per dose).

[0089] In some embodiments, the conjugated insulinotropic peptide can be administered as, for example, a single dose, a once-a-day dose, a once-a-week dose, a once-every-two weeks dose, a once-every-three-weeks dose or a once-a-month dose. In some embodiments, the weekly dose is administered once a week and the dose comprises the conjugated insulinotropic peptide in an amount between about 1000 to 4000 μg , preferably about 1250 to 3750 μg , and especially about 1250 to 2500 μg . Exemplary doses include 1000 μg , 1250 μg , 2500 μg and 3750 μg .

[0090] In certain aspects, the present invention provides unit dosages comprising a conjugated insulinotropic peptide, or a pharmaceutically acceptable salt thereof, in a form suitable for administration. Such forms are described in detail above. In certain embodiments, the unit dosage comprises 1000 to 4000 μg , 1250 to 3750 μg or 1250 to 2500 μg active ingredient. In particular embodiments, the unit dosages comprise about 1000, 1250, 1500, 1750, 2000, 2250, 2500, 2750, 3000, 3250, 3500, 3750 or 4000 μg active ingredient. Such unit dosages can be prepared according to techniques familiar to those of skill in the art.

[0091] Different therapeutically effective amounts of the conjugated insulinotropic peptide may be applicable for different disorders and conditions, as will be readily known by those of ordinary skill in the art. Similarly, amounts sufficient to prevent, manage, treat or ameliorate such disorders or conditions, but still sufficient to reduce nausea relative to administration of an unconjugated insulinotropic peptide are also encompassed by the above described dosage amounts and dose frequency schedules.

[0092] In certain embodiments, the methods and compositions can be practiced as a single, one time dose or chronically. By chronic it is meant that the methods and compositions of the invention are practiced more than once to a given individual. For example, chronic administration can be multiple doses of a pharmaceutical composition administered to a subject, on a weekly basis, monthly basis, or more or less frequently, as will be apparent to those of skill in the art. Chronic administration can continue for weeks, months, or years if appropriate according to the judgment of the practitioner of skill.

[0093] An effective amount of a conjugated insulintropic peptide described herein will provide therapeutic benefit with reduced nausea without causing substantial toxicity.

[0094] Toxicity of a conjugated insulintropic peptide can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, for example, by determining the LD50 (the dose lethal to 50% of the population) or the LD100 (the dose lethal to 100% of the population). The dose ratio between toxic and therapeutic effect is the therapeutic index. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a dosage range that is not toxic for use in human. The dosage of the compounds described herein lies preferably within a range of circulating concentrations that include the effective dose with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See, e.g., Fingl *et al.*, 1996, In: The Pharmacological Basis of Therapeutics, 9th ed., Chapter 2, p. 29, Elliot M. Ross).

5.3 Insulintropic Peptides

[0095] Useful insulintropic peptides include GLP-1, exendin-3 and exendin-4, and their precursors, derivatives and fragments. Such insulintropic peptides include those disclosed in U.S. Patent Nos. 6,514,500; 6,821,949; 6,887,849; 6,849,714; 6,329,336; 6,924,264; WO 03/103572 and 6,593,295, each of which are hereby incorporated by reference in their entirety.

[0096] In some embodiments, the insulintropic peptide is GLP-1. In some embodiments, the insulintropic peptide is a GLP-1 derivative. In some embodiments, the insulintropic peptide is exendin-3. In some embodiments, the insulintropic peptide is an exendin-3 derivative. In some embodiments, the insulintropic peptide is exendin-4. In some embodiments, the insulintropic peptide is an exendin-4 derivative. In some embodiments, the insulintropic peptide is exendin-4(1-39). In some embodiments, the insulintropic peptide is exendin-4(1-39)Lys⁴⁰.

5.3.1 GLP-1 and Its Derivatives

[0097] The hormone glucagon can be synthesized according to any method known to those of skill in the art. In some embodiments, it is synthesized as a high molecular weight precursor molecule which is subsequently proteolytically cleaved into three peptides: glucagon, GLP-1, and glucagon-like peptide 2 (GLP-2). GLP-1 has 37 amino acids in its unprocessed form as shown in SEQ ID NO: 1 (HDEFERHAEG TFTSDVSSYL

EGQAAKEFIA WLVKGRG). Unprocessed GLP-1 is essentially unable to mediate the induction of insulin biosynthesis. The unprocessed GLP-1 peptide is, however, naturally converted to a 31-amino acid long peptide (7-37 peptide) having amino acids 7-37 of GLP-1 ("GLP-1(7-37)") SEQ ID NO:2 (HAEG TFTSDVSSYL EGQAAKEFIA WLVKGRG). GLP-1(7-37) can also undergo additional processing by proteolytic removal of the C-terminal glycine to produce GLP-1(7-36) which also exists predominantly with the C-terminal residue, arginine, in amidated form as arginineamide, GLP-1(7-36) amide. This processing occurs in the intestine and to a much lesser extent in the pancreas, and results in a polypeptide with the insulinotropic activity of GLP-1(7-37).

[0098] A compound is said to have an "insulinotropic activity" if it is able to stimulate, or cause the stimulation of, the synthesis or expression of the hormone insulin. The hormonal activity of GLP-1(7-37) and GLP-1(7-36) appear to be specific for the pancreatic beta cells where it appears to induce the biosynthesis of insulin. Glucagon-like-peptide hormones are useful in the study of the pathogenesis of maturity onset diabetes mellitus, a condition characterized by hyperglycemia in which the dynamics of insulin secretion are abnormal. Moreover, glucagon-like peptides are useful in the therapy and treatment of this disease, and in the therapy and treatment of hyperglycemia.

[0099] Peptide moieties (fragments) can be chosen from the determined amino acid sequence of human GLP-1. The interchangeable terms "peptide fragment" and "peptide moiety" are meant to include both synthetic and naturally occurring amino acid sequences derivable from a naturally occurring amino acid sequence.

[00100] The amino acid sequence for GLP-1 has been reported by several researchers. See Lopez, L. C. *et al.*, 1983, *Proc. Natl. Acad. Sci.*, USA 80:5485-5489; Bell, G. I. *et al.*, 1983, *Nature* 302:716-718; Heinrich, G. *et al.*, 1984, *Endocrinol.* 115:2176-2181. The structure of the preproglucagon mRNA and its corresponding amino acid sequence is well known. The proteolytic processing of the precursor gene product, proglucagon, into glucagon and the two insulinotropic peptides has been characterized. As used herein, the notation of GLP-1(1-37) refers to a GLP-1 polypeptide having all amino acids from 1 (N-terminus) through 37 (C-terminus). Similarly, GLP-1(7-37) refers to a GLP-1 polypeptide having all amino acids from 7 (N-terminus) through 37 (C-terminus). Similarly, GLP-1(7-36) refers to a GLP-1 polypeptide having all amino acids from number 7 (N-terminus) through number 36 (C-terminus).

[00101] In one embodiment, GLP-1(7-36) and its peptide fragments are synthesized by conventional means as detailed below, such as by the well-known solid-phase peptide

synthesis described by Merrifield, J. M., 1962, *Chem. Soc.* 85:2149, and Stewart and Young, *Solid Phase Peptide Synthesis*, Freeman, San Francisco, 1969, pp. 27-66), which are incorporated by reference herein. However, it is also possible to obtain fragments of the proglucagon polypeptide, or of GLP-1, by fragmenting the naturally occurring amino acid sequence, using, for example, a proteolytic enzyme. Further, it is possible to obtain the desired fragments of the proglucagon peptide or of GLP-1 through the use of recombinant DNA technology, as disclosed by Maniatis, T., et al., *Molecular Biology: A Laboratory Manual*, Cold Spring Harbor, N.Y., 1982, which is hereby incorporated by reference.

[00102] Useful peptides for the methods described herein include those which are derivable from GLP-1 such as GLP-1(1-37) and GLP-1(7-36). A peptide is said to be "derivable from a naturally occurring amino acid sequence" if it can be obtained by fragmenting a naturally occurring sequence, or if it can be synthesized based upon a knowledge of the sequence of the naturally occurring amino acid sequence or of the genetic material (DNA or RNA) which encodes this sequence.

[00103] Also useful are those molecules which are said to be "derivatives" of GLP-1 such as GLP-1(1-37) and especially GLP-1(7-36). Such a "derivative" has the following characteristics: (1) it shares substantial homology with GLP-1 or a similarly sized fragment of GLP-1; (2) it is capable of functioning as an insulinotropic hormone; and (3) using at least one of the assays provided herein, the derivative has an insulinotropic activity of at least 1%, 5%, 10%, 25% 50%, 75%, 100%, or greater than 100% of the insulinotropic activity of GLP-1.

[00104] A derivative of GLP-1 is said to share "substantial homology" with GLP-1 if the amino acid sequences of the derivative is at least 80%, and more preferably at least 90%, and most preferably at least 95%, the same as that of GLP-1(1-37).

[00105] Useful derivatives also include GLP-1 fragments which, in addition to containing a sequence that is substantially homologous to that of a naturally occurring GLP-1 peptide may contain one or more additional amino acids at their amino and/or their carboxy termini, or internally within said sequence. Thus, useful derivatives include polypeptide fragments of GLP-1 that may contain one or more amino acids that may not be present in a naturally occurring GLP-1 sequence provided that such polypeptides have an insulinotropic activity of at least 1%, 5%, 10%, 25% 50%, 75%, 100%, or greater than 100% of the insulinotropic activity of GLP-1. The additional amino acids may be D-amino acids or L-amino acids or combinations thereof.

[00106] Useful GLP-1 fragments also include those which, although containing a sequence that is substantially homologous to that of a naturally occurring GLP-1 peptide, lack one or more additional amino acids at their amino and/or their carboxy termini that are naturally found on a GLP-1 peptide. Thus, useful polypeptide fragments of GLP-1 may lack one or more amino acids that are normally present in a naturally occurring GLP-1 sequence provided that such polypeptides have an insulinotropic activity of at least 1%, 5%, 10%, 25%, 50%, 75%, 100%, or greater than 100% of the insulinotropic activity of GLP-1. In certain embodiments, the polypeptide fragments lack one amino acid normally present in a naturally occurring GLP-1 sequence. In some embodiments, the polypeptide fragments lack two amino acids normally present in a naturally occurring GLP-1 sequence. In some embodiments, the polypeptide fragments lack three amino acids normally present in a naturally occurring GLP-1 sequence. In some embodiments, the polypeptide fragments lack four amino acids normally present in a naturally occurring GLP-1 sequence.

[00107] Also useful are obvious or trivial variants of the above-described fragments which have inconsequential amino acid substitutions (and thus have amino acid sequences which differ from that of the natural sequence) provided that such variants have an insulinotropic activity which is substantially identical to that of the above-described GLP-1 derivatives. Examples of obvious or trivial substitutions include the substitution of one basic residue for another (*i.e.* Arg for Lys), the substitution of one hydrophobic residue for another (*i.e.* Leu for Ile), or the substitution of one aromatic residue for another (*i.e.* Phe for Tyr), etc.

[00108] In addition to those GLP-1 derivatives with insulinotropic activity, GLP-1 derivatives which stimulate glucose uptake by cells but do not stimulate insulin expression or secretion are useful for the methods described herein. Such GLP-1 derivatives are described in U.S. Pat. No. 5,574,008, which is hereby incorporated by reference in its entirety.

[00109] GLP-1 derivatives which stimulate glucose uptake by cells but do not stimulate insulin expression or secretion which find use in the methods described herein include:

R¹-Ser-Tyr-Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-Val-Xaa-Gly-Arg-R² (SEQ ID NO:3)

wherein R¹ is selected from:

a) H₂N; b) H₂N-Ser; c) H₂N-Val-Ser; d) H₂N-Asp-Val-Ser; e) H₂N-Ser-Asp-Val-Ser (SEQ ID NO:4); f) H₂N-Thr-Ser-Asp-Val-Ser (SEQ ID NO:5); g) H₂N-Phe-Thr-Ser-Asp-Val-Ser (SEQ ID NO:6); h) H₂N-Thr-Phe-Thr-Ser-Asp-Val-Ser (SEQ ID NO:7); i) H₂N-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser (SEQ

ID NO:8); j) H₂N-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser (SEQ ID NO:9); and, k) H₂N-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser (SEQ ID NO:10); l) H₂N-His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser (SEQ ID NO:__); m) H₂N-His-D-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser (SEQ ID NO:__). In the peptide, Xaa is selected from Lys and Arg and R² is selected from NH₂, OH, Gly-NH₂, and Gly-OH.

These peptides are C-terminal GLP-1 fragments which do not have insulinotropic activity but which are nonetheless useful for treating diabetes and hyperglycemic conditions as described in U.S. Pat. No. 5,574,008, which is hereby incorporated by reference in its entirety.

5.3.2 Exendin-3 and Exendin-4 Peptides

[00110] The exendin-3 and exendin-4 peptide can be any exendin-3 or exendin-4 peptide known to those of skill in the art. Exendin-3 and exendin-4 are 39 amino acid peptides (differing at residues 2 and 3) which are approximately 53% homologous to GLP-1 and find use as insulinotropic agents.

[00111] The native exendin-3 sequence is HSDGTFTSDLSKQMEEEEAVRLFIEWLKNGG PSSGAPPPS (SEQ ID NO:11) and the exendin-4 sequence is HGEGTFTSDLSKQMEEEEAVRLFIEWLKNGGPSSGAPPPS (SEQ ID NO:12).

[00112] Also useful for the methods described herein are insulinotropic fragments of exendin-4 comprising the amino acid sequences: exendin-4(1-31) (SEQ ID NO:13) HGEGTFTSDLSKQMEEAVRLFIEWLKNGGPY and exendin-4(1-31) (SEQ ID NO:14) HGEGTFTSDLSKQMEEAVRLFIEWLKNGGY.

[00113] Also useful is the inhibitory fragment of native exendin-4 comprising the amino acid sequence: exendin-4(9-39) (SEQ ID NO:15) DLSKQMEEAVRLFIEWLKNGGPSSGAPPPS.

[00114] Other exemplary insulinotropic peptides are presented in SEQ ID NOS:16-22.

HDEFERHAEGTFTSDVSSYLEGQAAKEFIAWLVKGRK	SEQ ID NO: 16
HAEGTFTSDVSSYLEGQAAKEFIAWLVKGRK	SEQ ID NO: 17
HGEGTFTSDLSKQMEEAVRLFIEWLKNGGPSSGAPPPSK	SEQ ID NO: 18
HSDGTFTSDLSKQMEEAVRLFIEWLKNGGPSSGAPPPSK	SEQ ID NO: 19
HGEGTFTSDLSKEMEEVRLFIEWLKNGGPY	SEQ ID NO: 20
HGEGTFTSDLSKEMEEVRLFIEWLKNGGY	SEQ ID NO: 21
DLSKQMEEAVRLFIEWLKNGGPSSGPPPS	SEQ ID NO: 22

[00115] Useful peptides for the methods described herein include peptides which are derivable from the naturally occurring exendin-3 and exendin-4 peptides. A peptide is said to be "derivable from a naturally occurring amino acid sequence" if it can be obtained by fragmenting a naturally occurring sequence, or if it can be synthesized based upon a knowledge of the sequence of the naturally occurring amino acid sequence or of the genetic material (DNA or RNA) which encodes this sequence.

[00116] Useful molecules for the methods described herein also include those which are said to be "derivatives" of exendin-3 and exendin-4. Such a "derivative" has the following characteristics: (1) it shares substantial homology with exendin-3 or exendin-4 or a similarly sized fragment of exendin-3 or exendin-4; (2) it is capable of functioning as an insulinotropic hormone and (3) using at least one of the assays provided herein, the derivative has an insulinotropic activity of at least 1%, 5%, 10%, 25% 50%, 75%, 100%, or greater than 100% of the insulinotropic activity of either exendin-3 or exendin-4.

[00117] A derivative of exendin-3 and exendin-4 is said to share "substantial homology" with exendin-3 and exendin-4 if the amino acid sequences of the derivative is at least 80%, and more preferably at least 90%, and most preferably at least 95%, the same as that of either exendin-3 or 4 or a fragment of exendin-3 or 4 having the same number of amino acid residues as the derivative.

[00118] Useful derivatives also include exendin-3 or exendin-4 fragments which, in addition to containing a sequence that is substantially homologous to that of a naturally occurring exendin-3 or exendin-4 peptide may contain one or more additional amino acids at their amino and/or their carboxy termini, or internally within said sequence. Thus, useful derivatives include polypeptide fragments of exendin-3 or exendin-4 that may contain one or more amino acids that may not be present in a naturally occurring exendin-3 or exendin-4 sequences provided that such polypeptides have an insulinotropic activity of at least 1%, 5%, 10%, 25% 50%, 75%, 100%, or greater than 100% of the insulinotropic activity of either exendin-3 or exendin-4.

[00119] Similarly, useful derivatives include exendin-3 or exendin-4 fragments which, although containing a sequence that is substantially homologous to that of a naturally occurring exendin-3 or exendin-4 peptide may lack one or more additional amino acids at their amino and/or their carboxy termini that are naturally found on a exendin-3 or exendin-4 peptide. Thus, useful derivatives include polypeptide fragments of exendin-3 or exendin-4 that may lack one or more amino acids that are normally present in a naturally occurring

exendin-3 or exendin-4 sequence provided that such polypeptides have an insulintropic activity of at least 1%, 5%, 10%, 25%, 50%, 75%, 100%, or greater than 100% of the insulintropic activity of either exendin-3 or exendin-4.

[00120] Useful derivatives also include the obvious or trivial variants of the above-described fragments which have inconsequential amino acid substitutions (and thus have amino acid sequences which differ from that of the natural sequence) provided that such variants have an insulintropic activity which is substantially identical to that of the above-described exendin-3 or exendin-4 derivatives. Examples of obvious or trivial substitutions include the substitution of one basic residue for another (*i.e.* Arg for Lys), the substitution of one hydrophobic residue for another (*i.e.* Leu for Ile), or the substitution of one aromatic residue for another (*i.e.* Phe for Tyr), etc.

5.4 Conjugated insulintropic peptides

[00121] Useful conjugated insulintropic peptides for the methods described herein include insulintropic peptides and their derivatives conjugated to a macromolecule. In preferred embodiments, the insulintropic peptide is conjugated to albumin. Several methods can be used to link an insulintropic peptide to albumin. In certain embodiments, the insulintropic peptide is linked to albumin according to any technique known to those of skill in the art. In some embodiments, the insulintropic peptide is modified to include a reactive group which can react with available reactive functionalities on albumin to form covalent linkages.

[00122] The reactive group is chosen for its ability to form a stable covalent bond with albumin, for example, by reacting with one or more amino groups, hydroxyl groups, or thiol groups on the serum protein or peptide. Preferably, a reactive group reacts with only one amino group, hydroxyl group, or thiol group on albumin. Preferably, a reactive group reacts with a specific amino group, hydroxyl group, or thiol group on albumin. A useful conjugate of the methods described herein comprises a modified peptide, or a modified derivative thereof, which is covalently attached to albumin via a reaction of the reactive group with an amino group, hydroxyl group, or thiol group on albumin. Thus, a useful conjugate comprises a modified peptide, or a modified derivative thereof, in which the reactive group has formed a covalent bond to albumin.

[00123] To form covalent bonds with the functional group on a protein, one may use as a chemically reactive group a wide variety of active carboxyl groups, particularly esters. The carboxyl groups are usually converted into reactive intermediates such as N-hydroxysuccinimide (NHS) or maleimide that are susceptible to attack by amines, thiols and

hydroxyl functionalities on the protein. Introduction of NHS and maleimide reactive groups on the peptide can be performed by the use of bifunctional linking agents such as maleimide-benzoyl-succinimide (MBS) gamma-maleimido-butyryloxy succinimide ester (GMBS), dithiobis-N-hydroxy succinimido propionate (DTSP), succinimidyl 3-(2-pyridyldithio propionate) (SPDP), succinimidyl *trans*-4-(maleimidylmethyl) cyclohexane-1-carboxylate (SMCC), succinimidyl acetylthioacetate (SATA), benzophenone 4- maleimide, *N*-((2-pyridyldithio)ethyl)-4- azidosalicylamide (PEAS; AET). Such bifunctional linkers will activate either carboxy or amino group on the peptide based on the choice of protecting groups.

[00124] Alternatively the addition of maleimide to the peptide can be performed through the use of coupling agent such as N,N, dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide, hydrochloride (EDAC) and the likes to activate derivatives like maleimidopropionic acid, [2-[2-[2-maleimidopropionamido(ethoxy)ethoxy] acetic acid, and subsequently react with an amine on the peptide. Similar agents like DCC and EDAC could also be used to add derivatives like maleimidoalkyl amines to carboxy moieties on the peptide.

[00125] Primary amines are the principal targets for NHS esters. Accessible ϵ -amine groups present on the N-termini of proteins react with NHS esters. However, ϵ -amino groups on a protein may not be desirable or available for the NHS coupling. While five amino acids have nitrogen in their side chains, only the ϵ -amine of lysine reacts significantly with NHS esters. An amide bond can form when the NHS ester conjugation reaction reacts with primary amines releasing N-hydroxysuccinimide. These succinimidyl-containing reactive groups are herein referred to as succinimidyl groups.

[00126] In particular embodiments, the functional group on albumin is the single free thiol group located at amino acid residue 34 (Cys34) and the chemically reactive group is a maleimido-containing group such as MPA. MPA stands for maleimido propionic acid or maleimidopropionate. Such maleimido-containing groups are referred to herein as maleimido groups.

[00127] In some embodiments, albumin is covalently linked to a succinimidyl or maleimido group on the insulinotropic peptide. In some embodiments, an albumin amino, hydroxyl or thiol group is covalently linked to a succinimidyl or maleimido group on the insulinotropic peptide. In some embodiments, albumin cysteine 34 thiol is covalently linked

to a [2-[2-[2-maleimidopropionamido(ethoxy)ethoxy]acetamide linker on the epsilon amino of a lysine of the insulinotropic peptide.

[00128] In a specific embodiment, the reactive group is a single MPA reactive group attached to the peptide, optionally through a linking group, at a single defined amino acid and the MPA is covalently attached to albumin at substantially a single amino acid residue of albumin, preferably cysteine 34. In a preferred embodiment, the albumin is recombinant human albumin. In certain embodiments, the reactive group, preferably MPA, is attached to the peptide through one or more linking groups, preferably AEEA, AEA, or octanoic acid. In certain examples of embodiments in which the reactive group, preferably MPA, is attached to the peptide through more than one linking group, each linking group can be independently selected from the group consisting preferably of AEA ((2-amino) ethoxy acetic acid), AEEA ([2-(2-amino)ethoxy)]ethoxy acetic acid), and octanoic acid. In one embodiment, the reactive group, preferably MPA, is attached to the peptide via 1, 2, 3, 4, 5 or 6 AEEA linking groups which are arranged in tandem. In another embodiment, the reactive group, preferably MPA, is attached to the peptide via 1, 2, 3, 4, 5 or 6 octanoic acid linking groups which are arranged in tandem. In certain embodiments, a linking group can comprise, for example, a chain of 0-30 atoms, or 0-20 atoms, or 0-10 atoms. In certain embodiments, a linking group can consist of, for example, a chain of 0-30 atoms, or 0-20 atoms, or 0-10 atoms. Those atoms can be selected from the group consisting of, for example, C, N, O, S, P.

[00129] In certain embodiments, the reactive group can be attached to any residue of the insulinotropic peptide suitable for attachment of such a reactive group. The residue can be a terminal or internal residue of the peptide. In certain embodiments, the reactive group can be attached to the carboxy-terminus or amino-terminus of the peptide. In advantageous embodiments, the reactive group is attached to a single site of the peptide. This can be achieved using protecting groups known to those of skill in the art. In certain embodiments, a derivative of the insulinotropic peptide can comprise a residue incorporated for attachment of the reactive group. Useful residues for attachment include, but are not limited to, lysine, aspartate and glutamate residues. The residue can be incorporated internally or at a terminus of the peptide. In certain embodiments, the reactive group is attached to an internal lysine residue. In certain embodiments, the reactive group is attached to a terminal lysine residue. In certain embodiments, the reactive group is attached to an amino-terminal lysine residue. In certain embodiments, the reactive group is attached to a carboxy-terminal lysine residue, for instance, a lysine residue at the carboxy-terminus of GLP-1, GLP-1(7-37) or exendin-4.

[00130] The manner of modifying insulintropic peptides with a reactive group for conjugation to a macromolecule will vary widely, depending upon the nature of the various elements comprising the insulintropic peptide. The synthetic procedures will be selected so as to be simple, provide for high yields, and allow for a highly purified product. Normally, the chemically reactive group will be created at the last stage of insulintropic peptide synthesis, for example, with a carboxyl group, esterification to form an active ester. Specific methods for the production of modified insulintropic peptides are described in U.S. Patent Nos. 6, 329,336, 6,849,714 or 6,887,849, the contents of which are hereby incorporated by reference in their entirety.

[00131] The conjugated insulintropic peptides of the methods described herein may also be non-specifically conjugated to albumin. Bonds to amino groups will generally be employed, particularly with the formation of amide bonds for non-specific conjugation. To form such bonds, one may use as a chemically reactive group coupled to the insulintropic peptide a wide variety of active carboxyl groups, particularly esters. While a number of different hydroxyl groups may be employed in these linking agents, the most convenient would be N-hydroxysuccinimide (NHS) and N-hydroxy-sulfosuccinimide (sulfo-NHS). Other linking agents which may be utilized are described in U.S. Pat. No. 5,612,034, which is hereby incorporated by reference in its entirety.

5.4.1 Insulintropic Peptide Synthesis

[00132] Insulintropic peptides may be synthesized by standard methods of solid phase peptide chemistry known to those of ordinary skill in the art. For example, insulintropic peptides fragments may be synthesized by solid phase chemistry techniques following the procedures described by Steward and Young (Steward, J. M. and Young, J. D., 1984, Solid Phase Peptide Synthesis, 2nd Ed. (Pierce Chemical Company, Rockford, Ill.) using an Applied Biosystem synthesizer. Similarly, multiple fragments may be synthesized then linked together to form larger fragments. These synthetic peptide fragments can also be made with amino acid substitutions at specific locations. For solid phase peptide synthesis, a summary of the many techniques may be found in J. M. Stewart and J. D. Young, 1963, Solid Phase Peptide Synthesis. (W. H. Freeman Co., San Francisco), and J. Meienhofer, 1973, Hormonal Proteins and Peptides, vol. 2, p. 46, Academic Press, New York). For classical solution synthesis see G. Schroder and K. Lupke, The Peptides, Vol. 1, (Academic Press, New York). In some embodiments, synthesis of the insulintropic peptides is as described in U.S. Patent Nos. 6, 329,336, 6,849,714 or 6,887,849, the contents of which are hereby incorporated by reference in their entirety.

5.4.2 Conjugation

[00133] Preferably, the peptide and macromolecule (preferably albumin) of the conjugates of the invention are present in the conjugate in a 1:1 molar ratio, or an approximately 1:1 molar ratio. In particular embodiments, the peptide and macromolecule (preferably albumin) of the conjugates of the invention are present in the conjugate in a 1:1 molar ratio. In a preferred embodiment, the peptide and serum protein are present in the conjugate in a 1:1 molar ratio, or an approximately 1:1 molar ratio, and the peptide is attached to the reactive group, optionally through a linking group, at substantially only one site on the peptide and the reactive group is attached to the serum protein at substantially only one site on the serum protein. In a particular embodiment, the peptide and serum protein are present in the conjugate in a 1:1 molar ratio and the peptide is attached to the reactive group, optionally through a linking group, at substantially only one site on the peptide and the reactive group is attached to the serum protein at substantially only one site on the serum protein.

[00134] Preferably, the serum protein is albumin. Preferably, the single site of attachment of the reactive group to albumin is preferably the thiol group of cysteine 34 of albumin (e.g., via a maleimide linkage). In a specific embodiment, the reactive group is a single MPA reactive group attached to the peptide, optionally through a linking group, at a single defined amino acid and the maleimide is covalently attached to albumin at substantially a single amino acid residue of albumin, preferably cysteine 34.

[00135] In a preferred embodiment, a conjugate of the invention is formed by contacting a modified peptide comprising a maleimido group with a thiol-containing serum protein, preferably albumin, under conditions comprising a pH of between about 4.0 to 8.0, preferably between about 5.0 to 8.0 and more preferably between about 6.0 to 7.2, thereby preferably forming a stable thioether linkage which cannot be cleaved under physiological conditions. In certain preferred embodiments, the serum protein is recombinant human albumin.

[00136] In one embodiment, the modified peptide is amidated at its C-terminal end. In another embodiment, the modified peptide is not amidated at its C-terminal end. A conjugate of the invention can also comprise such an amidated peptide.


[00137] In a preferred embodiment, a single reactive group is covalently attached at a defined site of the modified peptide. In a preferred embodiment of the conjugate, a single reactive group is covalently attached at a defined site of the modified peptide and the reactive group is covalently attached to a single defined site of albumin, preferably to the thiol group

of amino acid residue Cys34 of albumin. Preferably, the reactive group of a modified peptide or conjugate of the invention comprises a maleimide group and forms peptide:albumin conjugates of approximately a 1:1 molar ratio. In certain embodiments, a 1:1 molar ratio of peptide to serum protein is preferred over higher ratios because a 1:1 molar ratio provides better biological activity than higher ratios (see *e.g.*, Stehle *et al.* 1997 *Anti-Cancer Drugs* 8:677-685, incorporated herein in its entirety).

[00138] In a preferred embodiment, the albumin is recombinant human albumin. Specific methods for the production of preformed peptide: albumin conjugates are described in U.S. Provisional Application No. 60/791,241, the contents of which is hereby incorporated by reference in its entirety. Specific methods for the purification of peptide: albumin conjugates are described in U.S. Patent Application Publication No. 2005/0267293, the contents of which is hereby incorporated by reference in its entirety.


[00139] Albumin can be provided from different sources, such as purified from blood sample(s) and expressed by recombinant techniques including recombinant yeast or bacteria. In some embodiments, the albumin is serum albumin. In some embodiments, the albumin is human serum albumin. In some embodiments, the albumin is bovine serum albumin. In some embodiments, the albumin is recombinant serum albumin. In some embodiments, the albumin is recombinant human serum albumin. In some embodiments, the albumin is RECOMBUMIN® (Delta Biotechnology Ltd., Nottingham, U.K.). RECOMBUMIN® is a recombinant human albumin (rHA) that is produced in vitro using recombinant yeast technology, in which genetically modified yeast (*Saccharomyces cerevisiae*) secrete soluble rHA which is subsequently harvested, purified and formulated for use as an excipient for the manufacture of biologics or a coating for medical devices. The main advantage of rHA over HSA is that it is expressed in yeast with no human derived products used in the manufacturing process. The use of such controlled production methods leads to a purer product with less structural heterogeneity. Previous studies have indicated that there is no significant difference between soluble rHA and HSA in terms of their biochemical characteristics, radiolabelling efficiency and biological behavior *in vitro* and *in vivo*. See Dodsworth *et al.*, 1996, *Biotechnol. Appl. Biochem.* 24: 171-176.

[00140] In certain embodiments, the conjugate is according to the following:



wherein X is S, O, or NH of an amino acid of said protein. In certain embodiments, said protein is albumin. In certain embodiments, said protein is albumin and X is S of Cys 34 of said albumin. Albumin of the conjugate can be any albumin as described above.

[00141] In certain embodiments, the conjugate is according to the following:



wherein X is S, O, or NH of an amino acid of said protein. In certain embodiments, said protein is albumin. In certain embodiments, said protein is albumin and X is S of Cys 34 of said albumin. Albumin of the conjugate can be any albumin as described above.

5.5 Monitoring the Presence of Conjugated Insulinotropic Peptides

[00142] The blood of the mammalian host may be monitored for the activity of the conjugated insulinotropic peptides and/or presence of the conjugated insulinotropic peptides using any method known to those of skill in the art. In certain embodiments, the conjugated insulinotropic peptides may be monitored using assays of insulinotropic activity. In other embodiments, the conjugated insulinotropic peptides may be monitored using HPLC-MS.

5.5.1 Assays of Insulinotropic Activity

[00143] The insulinotropic property of a conjugated insulinotropic peptide may be determined by providing that compound to animal cells, or injecting that compound into animals and monitoring the release of immunoreactive insulin (IRI) into the media or circulatory system of the animal, respectively. The presence of IRI is detected through the use of a radioimmunoassay which can specifically detect insulin.

[00144] Although any radioimmunoassay capable of detecting the presence of IRI may be employed, it is preferable to use a modification of the assay method of Albano *et al.*, 1972,

Acta Endocrinol. 70:487-509. In this modification, a phosphate/albumin buffer with a pH of 7.4 is employed. The incubation is prepared with the consecutive condition of 500 µl of phosphate buffer, 50 µl of perfusate sample or rat insulin standard in perfusate, 100 µl of anti-insulin antiserum (Wellcome Laboratories; 1:40,000 dilution), and 100 µl of [125 I] insulin, giving a total volume of 750 µl in a 10 x 75-mm disposable glass tube. After incubation for 2-3 days at 4° C., free insulin is separated from antibody-bound insulin by charcoal separation. The assay sensitivity is generally 1-2 µl U/ml. In order to measure the release of IRI into the cell culture medium of cells grown in tissue culture, one preferably incorporates radioactive label into proinsulin. Although any radioactive label capable of labeling a polypeptide can be used, it is preferable to use 3 H leucine in order to obtain labeling of proinsulin. Labeling can be done for any period of time sufficient to permit the formation of a detectably labeled pool of proinsulin molecules; however, it is preferable to incubate cells in the presence of radioactive label for a 60-minute time period. Although any cell line capable of expressing insulin can be used for determining whether a compound has an insulinotropic effect, it is preferable to use rat insulinoma cells, and especially RIN-38 rat insulinoma cells. Such cells can be grown in any suitable medium; however, it is preferable to use DME medium containing 0.1% BSA and 25 mM glucose.

[00145] The insulinotropic property of a conjugated insulinotropic peptide may also be determined by pancreatic infusion. The in situ isolated perfused rat pancreas preparation is a modification of the method of Penhos *et al.*, 1969, *Diabetes* 18:733-738. In accordance with such a method, fasted rats (preferably male Charles River strain albino rats), weighing 350-600 g, are anesthetized with an intraperitoneal injection of Amytal Sodium (Eli Lilly and Co., 160 ng/kg). Renal, adrenal, gastric, and lower colonic blood vessels are ligated. The entire intestine is resected except for about four cm of duodenum and the descending colon and rectum. Therefore, only a small part of the intestine is perfused, thus minimizing possible interference by enteric substances with insulinotropic immunoreactivity. The perfusate is preferably a modified Krebs-Ringer bicarbonate buffer with 4% dextran T70 and 0.2% bovine serum albumin (fraction V), and is preferably bubbled with 95% O₂ and 5% CO₂. A nonpulsatile flow, four-channel roller-bearing pump (Buchler polystatic, Buchler Instruments Division, Nuclear-Chicago Corp.) is preferably used, and a switch from one perfusate source to another is preferably accomplished by switching a three-way stopcock. The manner in which perfusion is performed, modified, and analyzed preferably follows the methods of

Weir *et al.*, 1974, *J. Clin. Investigat.* 54:1403-1412, which is hereby incorporated by reference.

5.5.2 HPLC-MS

[00146] HPLC coupled with mass spectrometry (MS) can be utilized to assay for the presence of conjugated insulintropic peptides as is well known to the skilled artisan. Typically two mobile phases are utilized: 0.1% TFA/water and 0.1% TFA/acetonitrile. Column temperatures can be varied as well as gradient conditions. Particular details are disclosed in U.S. Pat. No. 6,887,849 and 6,329,336 which are hereby incorporated by reference.

6. EXAMPLES

[00147] The invention is illustrated by the following examples which are not intended to be limiting in any way.

6.1 Example 1: Conditioned Taste Aversion Test (CTA)

[00148] One model for studying gastrointestinal malaise in response to administration of a test agent is the conditioned taste aversion test (Peeters and Broekkamp, 1994, *J. Steroid Biochem. Mol. Biol.*, 49:417-427). Conditioned taste aversion is a form of associative learning in which animals learn to avoid flavors of substances where initial ingestion of the substance was followed by illness. In conditioned taste aversion, animals learn to associate a flavor, or conditioned stimulus, with the aversive effects of an administered agent, or unconditioned stimulus. Following conditioned stimulus – unconditioned stimulus pairings, animals reduce their consumption of the conditioned stimulus, and/or show a preference for other tastes when later presented with the conditioned flavor. A conditioned taste aversion test was performed to determine gastrointestinal malaise following administration of unconjugated exendin-4(1-39) versus exendin-4(1-39) Lys⁴⁰ ϵ -AEEA-MPA conjugated with albumin (hereinafter called CJC-1134-PC in the following examples). The conditioned stimulus was 0.1% saccharin flavored water with high palatability. Lithium chloride was administered as a positive control for an unconditioned stimulus having an aversive gastrointestinal effect. See Riley and Baril, 1976, *Anim. Learn. Behav.* 4:1S-13S; Olivier *et al.*, 1999, *Pharmacol. Biochem. Behav.* 64:213-220.

[00149] Eighty male Sprague-Dawley rats of 6 weeks in age were divided into eight dosing groups (n=10/group). On the day of dosing, food and water were removed at 0900 h. At 1300 h, all rats were offered 0.1% saccharin (the conditioned stimulus) for four hours. At 1700 h, the saccharin solution was removed and test agents (unconditioned stimuli) were

administered accordingly: group 1: vehicle, 0.9% NaCl, subcutaneous injection (SC); group 2: LiCl, 80 mg/kg, intraperitoneal injection (IP); group 3: CJC-1134-PC, 0.0125 nmol/kg (0.9 µg/kg), SC; group 4: CJC-1134-PC, 0.5 nmol/kg (35 µg/kg), SC; group 5: CJC-1134-PC, 2 nmol/kg (140 µg/kg), SC; group 6: human serum albumin, 0.5 nmol/kg (32.5 µg/kg), SC; group 7: exendin-4, 0.5 nmol/kg (2 µg/kg), SC; and group 8: exendin-4, 0.5 nmol/kg (2 µg/kg), IP.

[00150] Following injection, all groups were given food ad libitum and 2 bottles of water. At 72 h and 96 h post dose, a choice of tap water or 0.1% saccharin solution (conditioned stimulus) was offered, and intake of both solutions was measured for 12 hours. Mean saccharin preference ratios, representing the percentage of saccharin solution consumed relative to total liquid intake, were determined for each group (FIGS. 1 and 2).

[00151] CJC-1134-PC induced a significantly lower taste aversion response compared to unconjugated exendin-4. While saccharin solution consumption in rats dosed with unconjugated exendin-4 (0.5 nmol/kg, SC) represented only ~ 40% of the total liquid intake for the 12 h period beginning 72 h post dose, saccharin solution consumption in rats receiving CJC-1134-PC (0.5 nmol/kg, SC) represented over 80% of the total liquid intake over the same period. Similar results were observed at 96 h post dose (FIG. 2). These results indicate a significant reduction in gastrointestinal malaise in response to CJC-1134-PC compared to unconjugated exendin-4. None of the three CJC-1134-PC doses resulted in conditioned taste aversion because there no statistically significant difference in saccharin preference ratios when compared to the internal negative vehicle control. On the contrary, exendin-4 decreased significantly the preference saccharine ratio as compared to the vehicle group aversion.

6.2 Example 2: Kaolin Intake

[00152] Another model for assessing gastrointestinal malaise in the rat is the consumption of non-nutritive substances following administration of a toxic agent. Pica, the ingestion of dirt or a non-nutritive clay such as kaolin, is a behavior only observed when rats are thought to be ill. As rats appear to lack an emetic reflex, kaolin ingestion has been used as an indirect indicator of nausea in response to a toxic agent. Accordingly, kaolin ingestion was measured in rats injected with unconjugated exendin-4 versus CJC-1134-PC to assess tolerability.

[00153] Ninety male Sprague-Dawley rats of 6 weeks in age were housed individually and given food and water ad libitum. Habituation to kaolin was achieved by offering fresh

blocks of kaolin daily for 8 days before dosing. Rats were then assigned to nine treatment groups (n=10/group). On the day of dosing (0900 h), rats were given clean cages with minimal bedding, and test agents were administered accordingly: group 1: vehicle, 0.9% NaCl, SC; group 2: LiCl, 80 mg/kg, IP; group 3: CJC-1134-PC, 0.125 nmol/kg (9 µg/kg), SC; group 4: CJC-1134-PC, 0.5 nmol/kg (35 µg/kg), SC; group 5: CJC-1134-PC, 2 nmol/kg (140 µg/kg), SC; group 6: human serum albumin, 0.5 nmol/kg (32.5 µg/kg), SC; group 7: exendin-4, 0.5 nmol/kg (2 µg/kg), SC; group 8: exendin-4, 0.5 nmol/kg (2 µg/kg), IP; and group 9: LiCl, 127 mg/kg, IP.

[00154] Immediately following injection, animals were offered a fresh kaolin block in addition to food and water. Kaolin intake was measured every hour for a four-hour period immediately after dosing, and also at 6, 8, and 24 hours post dose (FIG. 3).

[00155] Gastrointestinal malaise was immediate in rats receiving LiCl as indicated by significant kaolin intake after 1 h. Minimal kaolin intake was observed for the first 2 h in rats receiving both exendin-4 and CJC-1134-PC relative to 0.9% NaCl; however, intake dramatically increased by 3 h post dose in exendin-4 treated rats (0.5 nmol/kg SC). In CJC-1134-PC treated rats (0.5 nmol/kg SC), kaolin intake was only moderately increased after 6 h relative to 0.9% NaCl treated rats, thus indicating an increased tolerability of CJC-1134-PC compared to unconjugated exendin-4.

6.3 Example 3: Toxicity Studies

[00156] Toxicity of CJC-1134-PC was assessed in 7-day and 28-day studies utilizing both rat and primate subjects. CJC-1134-PC was administered daily by subcutaneous injection.

6.3.1 Rat Toxicity Study

[00157] For the 7-day study, Sprague Dawley rats were divided into four dosing groups (n=5/sex/group), and CJC-1134-PC was injected at 0, 1, 5 or 25 mg/kg/day SC. Immediately following injection, rats were housed individually, and food consumption and changes in body weights were monitored. Overall, no mortality or serious signs of toxicity were observed throughout the 7-day period. A dose dependent decrease in food consumption was observed at days 1-3 in rats dosed at 5 and 25 mg/kg/day. However, by day 4, food consumption was similar to that observed in rats receiving vehicle only (FIG. 4). Similar decreases in water consumption and body weight were observed in rats dosed at 5 and 25 mg/kg/day; however recovery was observed by days 3-4. (FIG. 5). There were no effects on clinical chemistry, hematology, urinalysis, or organ weights, or on macroscopic evaluations.

[00158] For the 28-day study, rats were divided into five dosing groups (n=10 or 15/sex/group), and CJC-1134-PC was injected at 0, 1, 5 or 25 mg/kg/day SC. Recombinant human albumin (rHA) was also dosed at 23 mg/kg as an additional control.

[00159] Following injections, a transient decrease in activity and fecal output was observed. A decrease in food consumption was observed at 25 mg/kg from days 1-7 (FIG. 6). However, by day 8, food consumption was comparable across all dosing groups. Decreases in body weight were also observed at the 25 mg/kg/doses at days 1-4. However, weight gain was comparable across all dosing groups from days 4-28 (FIG. 7). There were no effects on macroscopic evaluations. CJC-1134-PC was generally well tolerated in rats at up to 25 mg/kg.

6.3.2 Monkey Toxicity Study

[00160] For the 7-day study, monkeys were divided into 4 dosing groups (n=2/sex/group), and CJC-1134-PC was injected at 0, 0.4, 2, and 10 mg/kg/day SC. Overall, no mortality or serious signs of toxicity were observed throughout the 7-day period. Initial decreases in food consumption, as measured by the number of biscuits consumed per day, were observed across all dosing groups including vehicle control for days 1-3. By days 4-7, food consumption was comparable across all dosing groups (FIG. 8). There was a dose-dependent decrease in body weight by day 3 which was moderately attenuated by day 7 (FIG. 9). There were no effects on clinical chemistry, hematology, urinalysis, or organ weights, or on macroscopic evaluations.

[00161] For the 28-day study, monkeys were divided into 5 dosing groups (n=3 or 6/sex/group), and CJC-1134-PC was injected at 0, 0.4, 2, 10 mg/kg/day SC. Recombinant human albumin was also dosed at 9 mg/kg as an additional control. Food consumption was significantly reduced at 10 mg/kg on days 2 and 3 (FIG. 10). A transient decrease in activity, hunched posture, and decreased fecal output were also observed. However, food consumption was comparable across all dosing groups by day 7. A dose dependent decrease in body weight was observed throughout the 28-day period (FIG. 11). There were no effects on macroscopic evaluations. CJC-1134-PC was generally well tolerated in monkeys up to 10 mg/kg.

6.4 Example 4: Diabetes Test

[00162] The efficacy of CJC-1134-PC to treat diabetes was examined in 7 week-old female diabetic db/db mice (body weight range: 30.3 to 34.0 g; glycemia range: 8.6 to 21.9 mmol/L, pre-dose). Mice were divided into four groups and injected once by subcutaneous bolus injection (dose volume of 8 ml/kg) accordingly: group 1: vehicle, 0.9% NaCl, n=3;

group 2: CJC-1134-PC, 100 nmol/kg (7105 µg/kg), n=5; group 3: exendin-4, 100 nmol/kg (418 µg/kg), n=5; and group 4: exendin-4, 1 nmol/kg (4 µg/kg), n=5.

[00163] Blood glycemia was recorded at the following time points: pre-dose, 1, 3, 6, 24, 30, 48, 52, 74 and 101 hours post dose using a handheld glucometer (Accu-check compact, Roche Diagnostics, Canada). Food and water intake were recorded at the following timepoints: 0 to 6, 6 to 24, 24 to 30, 30 to 48, 48 to 72 and 72 to 101 hours post dose.

[00164] At 24 h post dose, blood glucose levels were significantly lower in mice receiving 100 nmol/kg CJC-1134-PC compared to 100 nmol/kg unconjugated exendin-4 (FIG. 12), and remained lower out to 48 h post dose (FIG. 13). After 72 h, blood glucose levels in CJC-1134-PC-treated mice were comparable to those observed in unconjugated exendin-4-treated mice (FIG. 14).

[00165] Food and water consumption was significantly reduced in mice dosed with CJC-1134-PC compared to unconjugated exendin-4 after 24 hours, and remained lower during the period of 24-47h post dose (FIG. 15 and FIG. 16). After 74 h, food consumption was only moderately lower in CJC-1134-PC treated mice relative to exendin-4 treated mice, and this difference was maintained out to 101 h post dose (FIG. 15). Water consumption remained moderately lower for CJC-1134-PC treated mice out to 101 h post dose (FIG. 16). No abnormal clinical signs were recorded during the study.

6.5 Example 5: Anti-Obesity tests

[00166] The effect of CJC-1134-PC and unconjugated exendin-4 on body weight gain and food intake was assessed in a diabetic obese rat model. Male Zucker Diabetic Fatty (ZDF) rats were divided into five groups and received four weekly subcutaneous injections accordingly: group 1: vehicle, 0.9% NaCl, n=8; group 2: exendin-4, 100 nmol/kg (418 µg/kg), n=8; group 3: exendin-4, 200 nmol/kg (836 µg/kg), n=8; group 4: CJC-1134-PC, 100 nmol/kg (7.1 mg/kg), n=8; and group 5: CJC-1134-PC, 200 nmol/kg (14.2 mg/kg), n=8.

[00167] Control animals were injected with 0.9% NaCl. Animals were weighed twice a week and food intake was measured daily for the first week and four times a week thereafter for a total of 25 days. Statistical analysis was performed using two-way Analysis of Variance (ANOVA) followed by the Tukey's pair-wise comparison test. Significance was considered when P value was less than 0.05.

[00168] Cumulative weight gain was significantly reduced in rats dosed with CJC-1134-PC compared to 0.9% NaCl (FIG. 17). Statistical significance was observed at 200 nmol/kg CJC-1134-PC beginning at day 4 ($p \leq 0.01$), and at 100 nmol/kg CJC-1134-PC

beginning at day 11 ($p \leq 0.05$); significance at both doses was maintained for the duration of the study. While cumulative weight gain was also reduced in unconjugated exendin-4 treated rats compared to 0.9% NaCl, this decrease was not significant compared to the 0.9% NaCl group.

[00169] Reduction in cumulative food intake was observed in both CJC-1134-PC treated as well as unconjugated exendin-4 treated rats relative to the 0.9% NaCl group. However, this decrease reached statistical significance in CJC-1134-PC treated rats after the second injection, on day 9, while statistical significance was not observed until after the third injection, on day 16, in exendin-4 treated rats (FIG. 18).

6.6 Example 6: Incidence of nausea and vomiting in type 2 diabetic patients: Exenatide versus CJC-1134-PC

6.6.1 Exenatide

[00170] Calara *et al.*, 2005, *Clin. Ther.* 27(2):210-214, conducted a randomized open-label crossover study examining the effect of injection site on bioavailability of exenatide (synthetic exendin-4). It was reported by the authors that three patients who received an inadvertent 10-fold overdose of exenatide (100 μg *i.e.*, 24 nmoles) experienced severe nausea and vomiting, with one patient experiencing severe hypoglycemia requiring aid. In patients receiving a normal 10 μg (2.4 nmoles) dose of exenatide, mild to moderate nausea was experienced by 36%, and vomiting by 21% of patients respectively.

[00171] Additionally, in three different Phase III clinical trials of exenatide, which enrolled a total of 1446 patients (480 patients receiving Exenatide 5 μg BID, 483 receiving Exenatide 10 μg BID, and 483 patients receiving placebo for 30 weeks), the most commonly reported treatment-emergent adverse events were nausea, at an incidence of 43.5%. See Iltz *et al.*, 2006, *Clin. Ther.* 28(5):652-665. The incidence of severe nausea associated with Exenatide therapy ranged from 2% to 5% in each of the 3 trials. Severe nausea generally occurred more frequently in the groups that received Exenatide 10 μg compared to those that received Exenatide 5 μg . Withdrawals attributed to nausea occurred in 2% to 4% of patients in the Exenatide arms across the 3 trials, again generally more frequently in the groups that received 10 μg compared to those that received 5 μg . Otherwise, reported nausea was generally mild to moderate and occurred with greater frequency during the first 8 weeks of therapy compared to the last 22 weeks.

[00172] Currently, exenatide is prescribed at 5 or 10 μg per dose twice a day.

6.6.2 CJC-1134-PC

[00173] A Phase I/II single escalating dose clinical study for the treatment of Type 2 diabetes using CJC-1134-PC was conducted to evaluate safety and tolerability, and as a secondary endpoint, the pharmacokinetic and pharmacodynamic (duration of activity after one injection based on mean glucose reductions) profile of CJC-1134-PC in patients with stable Type 2 diabetes.

[00174] Patients enrolled in the trial had HbA1c levels between 6.5% and 11%. Patients previously treated with oral anti-diabetic agents discontinued therapy at least 1 week prior to dosing. Six cohorts were dosed subcutaneously at 310, 620, 1250, 2500, 5000 and 3750 µg of CJC-1134-PC. Injections were performed with a 29 gauge needle. Each cohort consisted of 7 patients (6 active, 1 placebo). The mean glucose values at baseline of the cohorts (without placebo) were 15.6, 12.2, 12.1, 11.2, 9.6 and 15.3 mmol/L, respectively.

6.6.2.1 Safety and Tolerability

[00175] There were no safety or tolerability issues reported in the first four cohorts (310, 620, 1250 and 2500 µg dose), specifically, no nausea, no vomiting and no injection site reactions. At the 5000 microgram cohort, symptoms linked to an over stimulation of the GLP-1 receptors were observed as a result of the rapid decrease in blood glucose (manifested by headache, dizziness, and light-headedness without documented hypoglycaemia). In view of the positive activity demonstrated at the 1250 and 2500 microgram cohorts, 3750 micrograms of drug was dosed. In each of the 3750 and 5000 µg cohorts, there was one mild and transient case of short-term gastric stasis (observed post-lunch on day one); no anti-emetic medications were needed for either of these cases nor were any anti-emetics needed throughout the trial.

6.6.2.2 Pharmacokinetic Profile

[00176] The pharmacokinetic profile exhibited slow absorption and a prolonged exposure, with plasma drug levels rising for 4 to 6 days and then declining thereafter. For the 310 and 620 µg cohorts, the plasma concentration declined slowly for approximately 1 to 2 weeks following the plasma peak. Plasma concentrations were dose linear. Mean terminal half-life ranged from 164 hours to 274 hours and was not dose-dependent.

6.6.2.3 Efficacy

[00177] The efficacy parameters were based on the group mean and individual mean glucose values. Glucose was measured 6 times per day (fasting, 2-hour post-breakfast, pre-lunch, 2-hour post-lunch, pre-dinner and bedtime) during the first week and 3 times per day (fasting, 2-hour post-breakfast and bedtime) for the remaining 5 weeks of the study. In addition, a placebo group was constructed for data analysis by pooling the patients (one per

cohort) that received placebo during the study. Limited or no efficacy was observed in the first two cohorts (310 and 620 micrograms). At the third cohort (1250 micrograms), the mean glucose reduction from baseline for the first week was 19% (ranging from 15% to 24% with a mean reduction on day 7 of 20%), giving an average glucose value for the week of 9.8 mmol/L. This glucose reduction was significant against baseline ($p=0.0007$) and against placebo ($p=0.045$). Prolonged glucose lowering effect was observed for more than 2 weeks. The glucose reductions of the subsequent cohorts were of similar magnitude and duration and confirmed this response. At the 2500 microgram cohort, the average glucose value for the first week was reduced to 9.5 mmol/L; the value for the 5000 microgram cohort for the first week was reduced to 8.4 mmol/L. (The American Diabetes Association (ADA) has a recommended target of HbA1C < 7%; a mean daily glucose level of 9.5 mmol/L corresponds to an HbA1C level of 7%). Furthermore, weight reduction was observed during the first week: average loss of 0.3 kg at day 7 for the placebo group, average loss of 0.4 kg at day 7 for the first two non-active cohorts had an average loss of 0.9 kg at day 7 for the three following cohorts where activity was observed on glucose.

[00178] A comparison of the incidence of nausea and vomiting in Type 2 diabetic patients injected with a single subcutaneous injection of Exenatide versus CJC-1134-PC is represented in TABLE 1.

Table 1

Drug	N	Dose (μ g)	Dose (nmoles)	Nausea (%)	Vomiting (%)
Exenatide	25	10	2.4	36%	21%
	3	100	24	100%	100%
CJC-1134-PC	6	310	4.4	0	0
	6	620	8.7	0	0
	6	1250	17.6	0	0
	6	2500	35.2	0	0
	6	3750	52.8	17%	17%
	6	5000	70.4	50%	17%

[00179] All publications, patents and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were

specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

CLAIMS


What is claimed is:

1. A method of administering an insulinotropic peptide to a nausea-sensitive subject comprising administering to said nausea-sensitive subject the insulinotropic peptide conjugated to albumin, wherein the conjugated insulinotropic peptide reduces nausea compared to administration of the unconjugated insulinotropic peptide.
2. The method of claim 1 further comprising the step of selecting the nausea-sensitive subject prior to administering said insulinotropic peptide conjugated to albumin.
3. The method of claim 1 wherein said nausea sensitivity is a result of the administration of unconjugated insulinotropic peptide.
4. The method of claim 1 wherein said administering of the conjugated insulinotropic peptide prevents nausea in the subject.
5. The method of claim 1 wherein said administering of the conjugated insulinotropic peptide eliminates nausea in the subject.
6. The method of claim 1 wherein the amount of conjugated insulinotropic peptide administered is between 1000 and 4000 μg .
7. The method of claim 1 wherein the amount of conjugated insulinotropic peptide administered is between 1250 and 3750 μg .
8. The method of claim 1 wherein the amount of conjugated insulinotropic peptide administered is between 1250 and 2500 μg .
9. The method of claim 1 wherein the amount of the conjugated insulinotropic peptide administered is 1250 μg .
10. The method of claim 1 wherein the amount of the conjugated insulinotropic peptide administered is 2500 μg .
11. The method of claim 1 wherein the amount of the conjugated insulinotropic peptide administered is 3750 μg .

12. The method of claim 1 wherein said administering of the conjugated insulinotropic peptide is in an amount effective to treat a disease or disorder.
13. The method of claim 12 wherein the disease or disorder is obesity or diabetes.
14. The method of claim 13 wherein the disease or disorder is obesity.
15. The method of claim 13 wherein the disease or disorder is diabetes.
16. The method of claim 15 wherein the disease or disorder is type II diabetes.
17. The method of claim 15 wherein the amount is effective to reduce average daily blood glucose in the subject by at least 10 % over one week after a single administration.
18. The method of claim 1 wherein the subject is human.
19. The method of claim 1 wherein the conjugated insulinotropic peptide comprises albumin cysteine 34 thiol covalently linked to a [2-[2-[2-maleimidopropionamido(ethoxy)ethoxy]acetamide linker on the epsilon amino of a lysine of said insulinotropic peptide.
20. The method of claim 18 wherein said lysine has been added to the native insulinotropic peptide sequence.
21. The method of claim 18 wherein said lysine has been added to the carboxy terminus of said insulinotropic peptide.
22. The method of claim 20 wherein the insulinotropic peptide is selected from the group consisting of glucagon-like peptide 1, exendin-3, and exendin-4.
23. The method of claim 21 wherein the insulinotropic peptide is exendin-4(1-39).
24. The method of claim 1 wherein the albumin is human serum albumin.
25. The method of claim 1 wherein the albumin is recombinant serum albumin.
26. The method of claim 1 wherein the albumin is recombinant human serum albumin.
27. The method of claim 1 wherein the conjugated insulinotropic peptide comprises recombinant human serum albumin cysteine 34 thiol covalently linked to a [2-[2-[2


maleimidopropionamido(ethoxy)ethoxy]acetamide linker on the epsilon amino of the carboxy terminal lysine of exendin-4(1-39)Lys⁴⁰.

28. The method of claim 1 wherein the conjugated insulinotropic peptide is purified.
29. The method of claim 1, wherein said administering is oral, intranasal, intravascular, intraarterial, intramuscular, subcutaneous, transdermal or rectal.
30. A method of selecting a subject for treatment with an insulinotropic peptide conjugated to albumin, comprising the step of identifying a nausea sensitive subject having a disease or disorder treatable with the insulinotropic peptide, wherein the subject is selected for treatment when the subject is prone to experiencing or is experiencing nausea.
31. The method of claim 29, wherein said subject has experienced nausea as a result of the administration of unconjugated insulinotropic peptide.
32. A method of administering an insulinotropic peptide to a subject comprising (a) selecting a nausea sensitive subject having a disease or disorder treatable with the insulinotropic peptide; and (b) administering to the subject the insulinotropic peptide conjugated to albumin wherein the conjugated insulinotropic peptide reduces nausea compared to administering the unconjugated insulinotropic peptide.
33. A unit dose pharmaceutical comprising 1250 µg of a conjugate according to the following:




wherein X is S of an amino acid of human serum albumin.

34. A unit dose pharmaceutical comprising 2500 µg of a conjugate according to the following:



wherein X is S of an amino acid of human serum albumin.

35. A unit dose pharmaceutical comprising 3750 µg of a conjugate according to the following:



wherein X is S of an amino acid of human serum albumin.

36. A method for the treatment of diabetes or obesity in a subject, comprising

- a. selecting a nausea-sensitive subject in need of said treatment, and
- b. administering an insulintropic peptide linked to albumin to the selected subject,

wherein the administration of the insulintropic peptide linked to albumin provides a reduced nausea side effect compared to the administration of the insulintropic peptide alone.

37. The method of claim 35, wherein the disease is diabetes.

38. The method of claim 35, wherein the disease is obesity.

39. The method of claims 35, 36 or 37, wherein the insulintropic peptide is exendin-4(1-39)Lys⁴⁰.

40. The method of claim 38, wherein the insulintropic peptide is linked to recombinant albumin.

41. The method of claim 39, wherein a maleimido-containing group is coupled to the insulintropic peptide and has reacted with recombinant albumin.

1/18

Figure 1






Figure 2



ANOVA factorial, Fisher's post-hoc test
* p<0.05 vs. vehicle

Figure 3



4/18

Figure 4
7-Day Toxicity Study in Rats:
Food Consumption Male




Figure 5
7-Day Toxicity Study in Rats:
Body Weight Male






Figure 6
Food Consumption in Male Rats
(28-day Treatment & 2-week Recovery)




7/18

Figure 7


Delta Body Weight in Male Rats
(28-day Treatment & 2-week Recovery)






9/18

Figure 9
7-Day Toxicity Study in Monkeys:
Mean Body Weight Change
Male and Female



10/18

Figure 10
Food Consumption in Male Monkeys
(28-day Treatment & 2-week Recovery)



11/18

Figure 11

Delta Body Weight in Male Monkeys
(28-day Treatment & 2-week Recovery)






Figure 12
Blood glucose following a single SC
injection in female db/db mice



13/18

Figure 13

Blood glucose following a single
SC injection in female db/db mice



14/18

Figure 14

Blood glucose following a single
SC injection in female db/db mice





Figure 15

Food consumption



16/18

Figure 16




Figure 17

Effect of CJC-1134-PC and exendin-4 on body weight gain and food intake in diabetic obese rats: cumulative % weight gain versus Day 1.





Figure 18



INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2006/001835

A. CLASSIFICATION OF SUBJECT MATTER

IPC: *A61K 47/48* (2006.01), *A61K 38/26* (2006.01), *A61K 47/42* (2006.01), *A61P 1/08* (2006.01),
A61P 3/04 (2006.01), *A61P 3/10* (2006.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K 47/48 (2006.01)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used)

Canadian Patent database, United States Patent database, ESPACENET, Delphion, PubMed, Scopus (Keywords: insulinotropic, albumin, nausea, side effects, diabetes, obesity and related terms)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	CONJUCHEM press release "Final results of PC-DAC™:Exendin-4 phase I/II trial for type 2 diabetes confirms excellent tolerability, positive efficacy, and extended duration of activity" September 12, 2006 (Retrieved from the Internet at: http://conjuchem.hyphenhealth.com/news/PR%20EN%20Exendin-4%20Sept%2012%202006.pdf)	1-41
P,X	CONJUCHEM press release "Phase I/II preliminary results of PC-DAC™:Exendin-4 trial for type 2 diabetes demonstrate excellent tolerability and positive efficacy" April 26, 2006 (Retrieved from the Internet at: http://conjuchem.hyphenhealth.com/EN%20PR%2004-26-2006.pdf)	1-41
P,X	CONJUCHEM press release "ConjuChem to initiate phase I/II clinical study of PC-DAC™:Exendin-4 for type 2 diabetes" February 27, 2006 (Retrieved from the Internet at: http://conjuchem.hyphenhealth.com/news/20060227.shtml)	1-41
X	CONJUCHEM press release "ConjuChem provides clinical update on DAC™:GLP-1 and PC-DAC™:Exendin-4" September 30, 2005 (Retrieved from the Internet at: http://conjuchem.hyphenhealth.com/news/20050930.shtml)	1-41

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 April 2007 (03-04-2007)

Date of mailing of the international search report

10 April 2007 (10-04-2007)

Name and mailing address of the ISA/CA
Canadian Intellectual Property Office
Place du Portage I, C114 - 1st Floor, Box PCT
50 Victoria Street
Gatineau, Quebec K1A 0C9
Facsimile No.: 001-819-953-2476

Authorized officer

Wesley Sharman 819- 934-2326

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2006/001835

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CA2501421 (BRIDON et al.) November 23, 2000	1-26, 28-32 and 36-41
X	CA2550050 (HANSEN et al.) June 30, 2005	1-26, 28-32 and 36-41
X	CA2434237 (GLAESNER et al.) June 13, 2002	1-26, 28-32 and 36-41
L	NAUCK et al. "Glucagon-like peptide 1 and its derivatives in the treatment of diabetes" <i>Regulatory Peptides</i> , June 15, 2005, 128, 135-148.	1-22, 24-26, 28-32, 36-38, 40 and 41
L	CONJUCHEM press release "ConjuChem hits primary end-point in monotherapy phase II clinical trial with DACT TM :GLP-1" July 14, 2004 (Retrieved from the Internet at: http://www.glucagon.com/Conjuchem07.14.04%20PhII%20mono.pdf)	1-22, 24-26, 28-32, 36-38, 40 and 41
A	ADAMS et al. "Safety and utilization of blood components as therapeutic delivery systems" <i>Current Pharmaceutical Biotechnology</i> , 2003, 4, 275-282.	
A	KIEFFER et al. "The glucagon-like peptides" <i>Endocrine Reviews</i> , 1999, 20(6), 876-913.	
A	KIM et al. "Development and characterization of a glucagon-like peptide 1-albumin conjugate" <i>Diabetes</i> , 2003, 52, 751-759.	
A	BAGGIO et al. "A recombinant human glucagon-like protein (GLP)-1-albumin protein (Albugon) mimics peptidergic activation of GLP-1 receptor-dependent pathways coupled with satiety, gastrointestinal motility and glucose homeostasis" <i>Diabetes</i> , 2004, 53, 2492-2500.	

INTERNATIONAL SEARCH REPORTInternational application No.
PCT/CA2006/001835**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons :

1. ☒ Claim Nos. : 1-32 and 36-41
because they relate to subject matter not required to be searched by this Authority, namely :

Although claims 1-32 and 36-41 are directed to methods of medical treatment of the human/animal body (Rule 39.1(iv) of the PCT), a search has been carried out based on the alleged effects of the insulinotropic peptide-albumin conjugate in the treatment of diabetes and/or obesity.
2. ☐ Claim Nos. :
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :
3. ☐ Claim Nos. :
because they are dependant claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows :

(see extra sheet)

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos. :
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos. :

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2006/001835

Continuation of Box No. III

The claims of the present application are directed to a plurality of alleged inventions as follows:

Group A - Claims 1-32 and 36-41 are directed to methods of administering an insulinotropic peptide to nausea-sensitive subjects in order to treat a disease or disorder treatable with the insulinotropic peptide while eliminating the nausea side effect of such an administration by administering the insulinotropic peptide conjugated to albumin.

Group B - Claims 33-35 are directed to unit doses comprising specific doses of a specific insulinotropic peptide-albumin conjugate.

Insulinotropic peptides conjugated to albumin are known in the art. As such, there is no single inventive feature linking the use of these insulinotropic peptide-albumin conjugates in the treatment of nausea-sensitive subjects with unit doses comprising specific doses of a specific insulinotropic peptide-albumin conjugate. This is particularly the case as there is nothing in the present description suggesting that these specific doses play any role in reducing or eliminating the nausea side effect associated with the administration of unconjugated insulinotropic peptide.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2006/001835

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
CA2501421	23-11-2000	AT226593T T	15-11-2002
		AT252601T T	15-11-2003
		AT318835T T	15-03-2006
		AU754770B B2	21-11-2002
		AU761591B B2	05-06-2003
		AU764103B B2	07-08-2003
		AU765753B B2	25-09-2003
		AU4774800 A	05-12-2000
		AU4855500 A	05-12-2000
		AU5027100 A	05-12-2000
		AU5139300 A	05-12-2000
		AU2003246500 A1	16-02-2004
		BR0010750 A	26-02-2002
		BR0010757 A	19-02-2002
		CA2363712 A1	23-11-2000
		CA2372338 A1	23-11-2000
		CA2373252 A1	23-11-2000
		CA2373680 A1	23-11-2000
		CA2488348 A1	05-02-2004
		CA2499211 A1	23-11-2000
		CA2505617 A1	23-11-2000
		CN1191273C C	02-03-2005
		CN1351611 A	29-05-2002
		CN1698881 A	23-11-2005
		DE60000665D D1	28-11-2002
		DE60000665T T2	26-06-2003
		DE60006100D D1	27-11-2003
		DE60006100T T2	01-07-2004
		DE60026300D D1	27-04-2006
		DE60026300T T2	16-11-2006
		DK1105409T T3	03-07-2006
		DK1180121T T3	01-03-2004
		EA3922 B1	30-10-2003
		EP1105409 A2	13-06-2001
		EP1171582 A2	16-01-2002
		EP1179012 A1	13-02-2002
		EP1180121 A1	20-02-2002
		EP1264840 A1	11-12-2002
		EP1530588 A2	18-05-2005
		EP1591453 A1	02-11-2005
		EP1598365 A1	23-11-2005
		EP1623994 A2	08-02-2006
		ES2185595T T3	01-05-2003
		ES2209885T T3	01-07-2004
		ES2257298T T3	01-08-2006
		JP2002544287T T	24-12-2002
		JP2003500341T T	07-01-2003
		JP2003508350T T	04-03-2003
		JP2003527312T T	16-09-2003
		JP2005239736 A	08-09-2005
		JP2005255689 A	22-09-2005
		JP2005263807 A	29-09-2005
		JP2006151986 A	15-06-2006
		JP2006514607T T	11-05-2006
		NO20015584 A	03-01-2002
		PT1105409T T	31-07-2006
		PT1180121T T	31-03-2004
		SI1105409T T1	30-06-2006
		SI1180121T T1	30-04-2004
		US6329336 B1	11-12-2001
		US6514500 B1	04-02-2003
		US6593295 B2	15-07-2003
		US6821949 B2	23-11-2004
		US6849714 B1	01-02-2005
		US6887470 B1	03-05-2005
		US7090851 B1	15-08-2006

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2006/001835

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
CA2501421 (continued)	23-11-2000	US6887849 B2	03-05-2005
		US7090851 B1	15-08-2006
		US7144854 B1	05-12-2006
		US2004127398 A1	01-07-2004
		US2004138100 A1	15-07-2004
		US2004266673 A1	30-12-2004
		US2005070475 A1	31-03-2005
		US2005176641 A1	11-08-2005
		US2005176643 A1	11-08-2005
		US2005187159 A1	25-08-2005
		US2006009377 A1	12-01-2006
		US2006058235 A1	16-03-2006
		US2006135426 A1	22-06-2006
		US2006135428 A1	22-06-2006
		WO0069900 A2	23-11-2000
		WO0069902 A1	23-11-2000
		WO0069911 A1	23-11-2000
		WO0070665 A2	23-11-2000
		WO2004011498 A2	05-02-2004
		ZA200106676 A	19-07-2002
		ZA200109110 A	13-06-2002
CA2550050	30-06-2005	AU2004298425 A1	30-06-2005
		CN1893980 A	10-01-2007
		EP1696962 A2	06-09-2006
		KR20060109940 A	23-10-2006
		MXPA06006746 A	18-08-2006
		NO20063242 A	12-07-2006
		WO2005058958 A2	30-06-2005
CA2434237	13-06-2002	AU2689702 A	18-06-2002
		BR0116024 A	13-12-2005
		CN1483041 A	17-03-2004
		CZ20031602 A3	12-04-2006
		EA5584 B1	28-04-2005
		EP1355942 A2	29-10-2003
		EP1724284 A2	22-11-2006
		HR20030455 A2	31-08-2004
		HU0302529 A2	28-10-2003
		IL155812D D0	23-12-2003
		JP2004528014T T	16-09-2004
		MXPA03005036 A	05-09-2003
		NO20032565 A	01-08-2003
		NZ525577 A	27-05-2005